

This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

## Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + Refrain from automated querying Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

### **About Google Book Search**

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at http://books.google.com/

# Math 1809.11



# SCIENCE CENTER LIBRARY

FROM THE BEQUEST OF

HORACE APPLETON HAVEN, OF PORTSMOUTH, N. H.

(Class of 1848.)



. . . i

# **QUATERNIONS**

AS THE

# RESULT OF ALGEBRAIC OPERATIONS

BY


# ARTHUR LATHAM BAKER, Ph.D.

Head of Department of Mathematics, Manual Training High School, Brooklyn, N. Y.



#### NEW YORK

D. VAN NOSTRAND COMPANY 23 MURRAY AND 27 WARREN STREETS 1911 Math 1809.11



Copyright, 1911

BY

D. VAN NOSTRAND COMPANY

THE SCIENTIFIC PRESS.
ROBERT DRUMMOND AND COMPANY
BROOKLYN. N. Y.

## PREFACE

Beginners in the subject of Quaternions are generally bewildered by the arbitrary manner in which the subject is developed. They are forcibly introduced into a new domain where the familiar rules of combination of symbols are not valid. New magnitudes are arbitrarily assumed, subject to arbitrary laws. The reader finds the logic consistent and the results concordant with those of his previous courses, but he hardly knows why. He finds himself in a new country, but thoroughly and bewilderingly uncertain as to how he got there.

It is in the attempt to avoid this uncertain journey, to lead the student from the known to the unknown by familiar steps, by steps which require no arbitrary limitations of former laws, but merely their adaptation to new circumstances, that these class notes have grown into their present shape.

The backbone of the method of presentation is the use of a one-to-one correspondence between the mathematical concept and what I have ventured to call its idiographic symbol, that is, a symbol whose spatial properties are the same as the mathematical properties of the concept it symbolizes. From this similarity of properties there exists a one-to-one correspondence between the results of spatial operations upon the symbols and the corresponding mathematical operations upon the concept.

These idiographic symbols are strokes, spherical shells, and vectors, corresponding respectively to magnitudes having size, and correlated sense of opposition, scalars, and magnitudes having size, sense, and direction.

Spatial operations upon these symbols are used as suggestions for a one-to-one corresponding interpretation for for the mathematical concept.

These spatial operations are rational and logical and require no "standing loose for a time to logical accuracy." As they are rational and logical, so their interpretations are rational and logical, and the reader does not lose his sense of logical sequence. There is no "removal of barriers, of limitations, of conditions." Multiplication is the same from beginning to end, whether applied to scalars, vectors, or quaternions. Commutivity of factors may be permissible in some cases and not in others, but this is a mere incident and not an essential element of the operation.

The reader is not mystified by arbitrarily defining multiplication of one vector into another as the turning through a right angle, etc., and left to wonder how one line can do anything to another. In fact, the operations are not defined a priori at all, but taking the properties of discrete quantities as symbols of operations which the reader is to perform, we find six possible operations, addition, subtraction, multiplication, division, reversion, and mean reversion. These operations are defined accordingly a posteriori as results of causes, not arbitrarily as assumptions.

The performance of these operations upon scalars leads to or *evolves* successively vectors and quaternions. Thus quaternions are evolved from discrete magnitudes, not arbitrarily, but of necessity, and along certain fixed and preordained

<sup>\*</sup> Kelland and Tait, and others.

lines, by rules which the properties of discrete magnitudes necessitate, and which cannot be altered or varied, and with which the reader is already familiar. The reader is not disturbed by the thought, Suppose we had made some other assumption, what then? No assumptions are made. He simply follows the road suggested by the properties of discrete magnitudes, and can arrive at but one result.

We make no laws, lay down no rules, make no modifications or limitations. The only way in which we exercise any choice is in the rational application of the laws we discover to the proper operands and in a proper and logical manner.

The "interpretation of our results" is not made to "depend upon the definition" as a foundation. The foundation is the properties of discrete magnitudes, and the definitions are merely rational statements of the results of these properties being used as suggestions for operations to be performed by the reader.

Considerable stress has been laid upon the avoidance of the sole use of mere typographical symbols and upon the auxiliary use of idiographic symbols upon which spatial operations can be performed; as in the use of strokes for merely reversible magnitudes, a spherical shell for scalar magnitudes, are strokes for quaternion multiplication, two vector factors for the corresponding quaternion, etc., thus making the treatment concrete and avoiding the difficulties of abstractness.

The original features of the book are those specified above, coupled with the general heuristic method by which the student hews out his own concepts as he goes along. The results, the examples, the applications, and the terms used are those found in every treatise on the subject, of which I have made free use and to whom should be accredited

these features: particularly Hamilton, Kelland and Tait, Tait, Laisant, Molenbrock, Hathaway.

As these notes are only intended as an introduction, not an overabundance of examples or formulæ has been provided, nor have any applications been made to problems in Geometry and Physics. These will be found in the works cited. Nor has the subject of differentiation been touched upon.

The author's own experience with this method of presentation of the subject to beginners has been encouraging. It is hoped others will have the same experience.

# CONTENTS

#### CHAPTER I

CHAPTER I	
1	PAGE
MATHEMATICAL OPERATIONS UPON DISCRETE MAGNITUDES	1
Discrete magnitudes, 1; Addition, 4; reversion, 5; subtraction, 6; multiplication, 7; division, 8; mean reversion, 9.	
CHAPTER II	
Idiographs	6
Strokes, 11; Argand diagram, 13; addition of strokes, 15; algebraic operations, 17; scalars, 18.	
CHAPTER III	
Space Idiographs	10
Space idiographs, 19; mean reversion, 20; vector, 22; reversal, 26; addition and subtraction of vectors, 27; decomposition of vectors, 29; notation, 30; exercises in combination, 31; examples, 37.	
CHAPTER IV	
MULTIPLICATION OF UNIT VECTORS	17
Parallel vectors, 40; perpendicular vectors, 41; inclined vectors, 42; exercises in i, j, k, 44.	
vii	

## CONTENTS

## CHAPTER V

PAGE

QUATERNIONS	21
Quaternion, 45; meaning of $\frac{1}{\alpha}$ , 47; examples, 48; commu-	
tivity of vectors, 49; types of quaternions, 50; notation, 53; exercises, 54; geometric meaning of $TV\alpha\beta$ and $S\alpha\beta$ , 58; meaning of $S\alpha\beta\gamma$ , 59; sum of scalar and vector, 61; definitions of algebra, etc., 63.	
CHAPTER VI	
Kinds of Quaternions	32
CHAPTER VII	
Quaternion Operators	37
CHAPTER VIII	
Products of Quaternions	45
CHAPTER IX	
Versors	48
Versors, 94; multiplication of versors, 97; examples, 99; conversion of $qr$ into $rq$ , 101; meaning of $qrq^{-1}$ , 102; examples, 103.	

ix

# CHAPTER X PAGE Interpretation of Vector Equations..... 54 $\alpha\beta\gamma=8$ , 107; exercises and formulæ, 111; applications. 114; reference formulæ, 122. CHAPTER XI QUATERNION EQUATION OF THE FIRST DEGREE ..... General form, 127; linear scalar equation, 129; decomposition of vector part, 130; linear vector function, 131; properties of $\phi$ , 132; conjugate strain function, 135; application of $\phi$ to vector, 141; properties of $\phi$ , 143; nonion, 145. CHAPTER XII 71 Change of volume, 146; special values of modulus, 148; special applications of $\phi$ , 153; properties of $(\phi'-g)$ , 154; latent lines, latent roots, latent planes, 162; two latent roots equal, 164; three latent roots equal, 165; pure strain, 172; examples, 175. APPENDIX.....

Functional Symbols.

.

# **QUATERNIONS**

### CHAPTER I

# MATHEMATICAL OPERATIONS UPON DISCRETE MAGNITUDES

1. Mathematics is for school purposes the science of magnitude.

Magnitude has been defined as that which can be increased or diminished, and of two kinds, continuous magnitude (continuum), which is usually called quantity and answers to the question, How much? and discrete magnitude (discreta), which is usually called number and answers to the question, How many?

Quantity is differentness from nothingness, or quantitatively expressed, difference from nothing.

2. When we attempt to express quantity in symbols or terms, or to conceptualize it, we find ourselves compelled to express it in terms of some arbitrary unit adopted as a standard. The symbol or concept for the measure of quantity is called *number*, expressing its differentness from oneness, the oneness of the standard.

In the case of natural units, soul, personality, inhabitant, etc., the differentness from oneness which distinguishes a group from an individual is number, giving rise to the so-called primary numbers, 1, 2, 3, ...

Number is differentness from oneness. The whole

essence of number is to differ from another number. It belongs to the primary categories of time, space, matter, etc. No one of these can exist without the presence of number, that is without a oneness and a difference from that oneness. The very existence of time, space, matter implies multiplicity, three-foldness.

The limitations of the mind require symbols to represent the different concepts so that the mind can posit its judgments and conclusions to await its return to make use of them. Otherwise the mind soon becomes clogged with the impedimenta of its own creation.

Upon the fortunate choice of these symbols depends very greatly the progress in their use. A striking illustration of this is seen in the enormous advantage of the Hindu position system over the clumsy notation of the Greeks and Romans.

3. In the infancy of the subject pebbles (calculi) and other concrete objects were the symbols used. And as a negative pebble was impossible of imagination, negative numbers were imaginary. Even as late as the sixteenth century they were called by Cardan in his Ars Magna, numeri ficti, imaginary numbers. After they were dignified with the title of real numbers, expressions containing  $\sqrt{-1}$  would intrude themselves, under the operations of mathematical analysis, and as there was no way of writing a numerical symbol whose square would be a negative number, these were in their turn imaginaries, even to the present day. So late a writer as De Morgan (1831) speaks of them as void of meaning, self-contradictory, and absurd, though of great utility in the formal mechanism of Algebra.

But it is not the number that is imaginary, it is the application of it to certain data or symbols that produces the imaginary features. Fractions are imaginary when counting souls, inhabitants, events, phenomena, etc. Negative

numbers are imaginary when applied to length, bulk, intensity, etc. The imaginaries of to-day become real when applied to the proper symbol.

Before searching for a symbol we must inquire about the properties of primary number and the operations performable upon it. What are these performable operations, guiding ourselves by the properties of number itself? This guidance is necessary, the properties of a thing, if we thus speak of number, being the determining factors as to what purposes it can be put.

- 4. Number is differentness from unity. We can increase this differentness quantitatively by combining the differentness of one number with the differentness of another number. This operation we call addition, and by means of it we can evolve all the primary numbers from unity. In this two numbers are taken accumulatively.
- 5. We can imagine the differentness from oneness to have a correlated sense of opposition, like debit and credit, affirmation and negation, an antagonistic differentness. One is the reversal of the other, the negative of the other; and the process of converting one into the other is called reversion.

Numbers having this correlated sense of opposition are said to have size and sense. An example could be the accumulative collection of material into the form of a mound or tumulus. The workmen could proceed decumulatively (this word is not in the lexicons, but its usefulness is apparent) by undoing their former work, and might proceed below the bed rock. We might not call this a negative tumulus, but it would be the negative of the tumulus concept and the number concept would be a negative number.

6. We can diminish the differentness from oneness which is characteristic of a number by the characteristic different-

ness of some other number, giving rise to the operation called subtraction. The continued application of this process to numbers having size and sense leads to negative numbers. Here the numbers are taken decumulatively. As addition is accumulative combination so subtraction is decumulative combination of numbers without regard to the size of the subtrahend.

We can consider the characteristic differentness of one number as a mandate for an operation to be performed upon another number. Now this characteristic differentness from oneness or unity may be evolutory or involutory, differentness from unity, or differentness toward unity, a differentness which evolves the number from unity, or a differentness which converts a number into unity.

7. If we take the evolutory differentness as the mandate, then the operation is called multiplication, the doing to the operand (multiplicand) what was done to unity to produce the operator (multiplier). If the first operand and all the operators in a series of operations are the same, the operation is called involution. Involution works from unity with the first operand and all the operators alike.

A curious error is liable to creep in here unless care be exercised. For example, in  $\sqrt{2}.3$ , one is liable to say that  $\sqrt{2}$  is derived from 1 by doubling 1 and taking the  $\sqrt{\phantom{0}}$  of the result. This applied to 3 gives the result  $\sqrt{2.3}$ , which is, of course, wrong. The error consists in taking the square root of the result, whereas multiplication is the doing what was done to unity, and not what was done to the results of operations upon unity. To get  $\sqrt{2}$  from unity, we really operate upon unity with the tensor or stretcher  $1.4142...=\sqrt{2}$ .

8. If we take the differentness toward unity as the mandate, then the operation is called division, the doing to the

operand (dividend) what was done to the operator (divisor) to produce unity.

If all the operators and the last operand in a series of operations are the same, the operation is called **evolution**, and gives rise to irrational and so-called imaginary numbers, e.g.,  $\sqrt{2}$ ,  $\sqrt{-1}$ , etc.

Evolution works towards unity with all the operators and the last operand alike.

**9.** One of the most important phases of evolution is  $\sqrt{a}$ , the breaking up of the operation of passing from unity to a into two equal steps, so that the repetition of the first step,  $\sqrt{a}$ , shall produce a. When a=-1, we get  $\sqrt{-1}$ .

Since -1 is reversion,  $\sqrt{-1}$  is called mean reversion.

Unlike addition and subtraction, which merely adjoin the elements without change, multiplication and division are transformational operations which change the operand into an entirely new and different number. They are purely algebraic, being limited, unlike addition and subtraction, to operations upon discrete number. A not entirely satisfactory illustration of the difference between the two operations of addition and multiplication is the putting of two canes together for addition; for multiplication we would be compelled because the first was triangular, black, smooth, and uniform in size to transform the second cane which was round, white, knotted, tapering, and with a knob on the end, into a triangular, black, smooth, prismatic It is transformed into an entirely new and distinct form, and its original features have disappeared. numbers, addition and subtraction are merely adjunction without change of the original elements. Multiplication and division are destructive transformations, the destruction of one element and the production of an entirely new and different one, like the growth of a plant from a seed.

#### CHAPTER II

#### IDIOGRAPHS

10. Mathematical operations are conducted by means of symbols, and upon our choice of symbols depends largely the success of our operations.

In the dawn of mathematics, the fingers or other concrete objects were the symbols used, and a negative quantity was purely imaginary. With the introduction of written symbols of quantity and the concept of debit and credit, the negative quantity lost its imaginary quality and became real. In these symbols, however,  $\sqrt{-a}$  remained imaginary, because the spatial and typographical properties of the symbol and the mathematical properties of the thing it represented did not agree.

11. With the introduction of the symbol called a stroke, a straight line upon the surface of the paper, to represent a magnitude of a given size, the space properties of the symbol and the mathematical properties of the thing symbolized became the same, and spatial operations upon the symbols corresponded to mathematical operations upon the thing symbolized, and could be used to interpret and control the mathematical operations.

Symbols whose spatial properties are the same as the mathematical properties of the things represented might be called *idiographs* (ĩδως, proper, peculiar).

12. Representing a magnitude having sense of generation, sense of correlated opposition, by the idiograph ——, and

of course its reversal by  $\leftarrow$ —, how can we convert —— into  $\leftarrow$ —?


Obviously, only in one way, so long as we retain its stroke characteristics, namely, by swinging it through an angle of 180° in the surface of the paper. To attempt to swing it out of the surface of the paper is to lose its stroke characteristics, to give it absolute direction in space and render it no longer an idiograph.

The revolution of 180° can be broken into two equal steps of 90° each. Hence  $\uparrow$  or  $\downarrow$  must be the idiographic equivalent of  $\sqrt{-}$ .

Combining these into one diagram and assuming the normal revolution as counter-clockwise, we get the idiographic diagram with the corresponding typographical symbols.



13. This is the well-known Argand diagram, affording a simple method of representing relatively directed quantities, or as they are generally called, *complex quantities*, the general type of which is  $x+\sqrt{-1}y$ , where x represents the normal or the reversed portion and  $\sqrt{-1}y$  the mean reversed




portion. It is generally written x+iy, i standing for  $\sqrt{-1}$ . The rectangular co-ordinates x and iy determine a point, whose distance from the origin, r, is called the **modulus**, and whose angular distance

from the axis of x, the angle  $\phi$ , is called the **amplitude** of the point or complex quantity x+iy.

14. The idiographic symbol for a magnitude, —, showing its size and sense, we have already designated as a stroke, a stroke forward or a stroke backward. Two forward strokes need not be represented by the same

symbol on the paper. The forwardness is in reference to its own backwardness, and has no reference to the forwardness or backwardness of other strokes. A stroke is a straight line in a plane, symbolizing a given magnitude in size and in relation to its sense of normalcy or of reversion, or a condition between these. Two strokes are equal when they have the same lengths and the same direction in a plane as regards a standard normal direction in the plane.

15. If several strokes be taken in succession the sum (result) of them is the same as the stroke from the beginning of the first, to the end of the last, when they are arranged end to end so as to be successive. Thus



$$\alpha + \beta + \gamma + \delta = \varepsilon$$
.

Strokes will for the present be represented typographically by lowercase Greek letters, as above.

The reader must notice carefully that it is not the lengths of the stroke which are added, but the results of

the strokes, including both length and direction on the paper.

It is easily seen that the order of the strokes is immaterial, and that any number of consecutive strokes can be replaced by their sum. The addition of strokes is a commutative and associative operation, that is, the order and mode of grouping has no effect on the result.

A stroke is subtracted by reversing its direction and adding.

16. If we attempt to break up the operation of reversal into three or more equal and similar operations, for example, three, as shown in the diagram, we find

that  $\sqrt[3]{-1} = \frac{1}{2} + i\frac{\sqrt{3}}{2}$ , that is, the first of the three equal operations is expressible in terms of the operation of mean reversion, and of course the second operation likewise.

Similarly for a greater number of steps. Hence mean reversion is evidently the unique operation, a multiple of which is reversion, all the other partial equal operations whose continued application results in reversion being expressible in this one.

17. Hence there are six unique and fundamental operations which can be performed upon a discrete magnitude: addition, subtraction, reversal, multiplication, division, and mean reversion, and no others.

By fundamental operations is meant operations based upon the properties of discrete magnitudes, size, and sense of correlated opposition.

18. Mere discrete magnitudes, considering size only, are scalars, that is, they can be scaled off on a scale either in a normal or in a reversed direction.

#### CHAPTER III

#### SPACE IDIOGRAPHS

19. Space idiographs. In space we cannot idiographically represent a scalar by a line, for that would be assigning to the symbol a characteristic direction, which the magnitude it represents does not possess. If we are to represent a scalar magnitude in space by any idiographic symbol, the only one which seems available as possessing perfect symmetry and therefore devoid of direction is a spherical shell.

Just as we can assume our + unit of heat, pressure, etc., anywhere on the scale, so we can posit our +spherical shell anywhere in space.

Likewise as the — unit of heat, etc., would naturally adjoin the + unit, along the scale, so naturally we should expect the — unit shell to adjoin the + unit shell in some position determined by previous assignment of direction. According to this previously determined direction we shall have units of  $\alpha$  direction, of  $\beta$  direction, etc., where  $\alpha$  and  $\beta$  denote direction, not magnitudes, unless we say unit magnitude, just as previously we had units of heat sense, credit sense, etc.

20. How can we break up the operation of transforming the + shell into the - shell into two similar and equal operations. The + shell can be changed into the - shell by the repetition of two different operations:


A. By revolving the — shell about its point of contact with the + shell through an angle of 90° twice;

- **B.** By moving the elements of the shell perpendicularly to the common line of centers in the proportion  $\sin \theta$  (where  $\theta = \cos^{-1}$  harmonic displacement \* of the shell element) and moving the resulting configuration, a directed line, one-half its dimensions toward the correlatively reversed position of the original operand, i.e., toward the position of the + shell. A repetition of this operation would produce the + shell.
- 21. Operation A is excluded by reason of its lack of definiteness, leaving operation B as the operation producing the mean reversed state.

Since we are dealing with iodiographs these spatial operations must have a one-to-one correspondence with mathematical operations performed upon the things they symbolize.

- 22. Hence a mean reversed scalar is represented in all its properties by a directed magnitude in space, a vector, as it is called, which is definitely directed as soon as the + and shells are posited.
- 23. A vector is any magnitude having direction of extension in space, a directed line, plane, etc., such as velocity, impulse, force, etc.

Vectors are equal when they possess the same quantitative



\* If P represents an element of the shell, OA, the projection of the radius on a given diameter is its harmonic displacement,

 $\theta = \cos^{-1} OA$ .


and qualitative properties, viz., magnitude, sense, and direction of extension. Direction of extension is that property which prevents the vectors from coinciding (in whole or in part) when brought together.

Parallel vectors of the same length are equal. Vectors can be made coinitial without altering their properties.

- **24.** It is customary to indicate the unit vector in a given direction by Greek letters  $\alpha$ ,  $\beta$ ,  $\gamma$ , ... Generally three unit vectors at right angles to each other are assumed as reference units. These are designated by i, j, k.
- 25. So far we have recognized two kinds of magnitudes, scalars and vectors, and six operations, addition, subtraction, reversion, multiplication, division, and mean reversion.

Applying these operations to scalars we find that they all produce scalars again, except in the case of mean reversion, and that produces a vector. This gave us the second kind of magnitude, to which we will now proceed to apply the six fundamental operations.

- 26. Reversal is merely the turning of the vector into the opposite direction, as the word implies. The result is sometimes called a revector.
- 27. Addition and subtraction of vectors. Subtraction of vectors is merely addition with the minuend reverted. Vectors are of the nature of strokes, with the property of absolute direction added. The laws governing the addition



of strokes evidently hold here also. Thus, vector addition is commutative and associative, and this whether the vectors are coplanar or not. Thus

$$\alpha + \beta + \gamma = \beta' + \gamma' + \alpha'$$

$$= \gamma'' + \alpha'' + \beta'', \text{etc.},$$

where  $\alpha$ ,  $\beta$ ,  $\gamma$  are the three edges of a parallelopiped. The same reasoning would apply to additional vectors.

28. The following equations are self evident:

$$\alpha + \alpha + \alpha$$
 ... to  $m$  terms  $= m\alpha$ ,  
 $-\alpha + (-\alpha) + \ldots$  to  $m$  terms  $= m(-\alpha) = -m\alpha$ ,  
 $m\alpha + m\beta = m(\alpha + \beta)$ ,  $m\alpha - m\beta = m(\alpha - \beta)$ .

**29.** If  $\alpha$ ,  $\beta$ ,  $\gamma$  be three coinitial vectors, then any fourth coinitial vector,  $\xi$ , can be expressed as

$$\xi = x\alpha + y\beta + z\gamma$$

 $\xi$  being the diagonal of the parallelopiped whose edges are  $x\alpha$ ,  $y\beta$ , and  $z\gamma$ .

**30.** If  $\alpha$  is a unit vector, and  $m\alpha = A$ , then m indicated generally by the symbol TA, which expresses the length of the vector A, is called the **tensor** (tendere, to stretch) of the vector A.  $\alpha$ , denoted by UA is called the **unit vector** of A. Therefore

$$A = TA \cdot UA$$

Vectors will be denoted by capital Greek letters when the tensor and unit part are to be emphasized; by lower-case Greek letters when the question of length is not important; and by the corresponding lower-case English and Greek letters when speaking of the tensor and unit part separately.

Thus the same vector may be indicated by A,  $a\alpha$ ,  $\alpha$ .


The tensor is signless, just as any length, the height of a steeple, for instance, is signless; or the height of a man. A man cannot be -5 feet tall.

### EXERCISES IN VECTOR COMBINATIONS

31. If  $l\alpha + m\beta = 0$ , then l = 0, m = 0, for in no other way can two strokes in different directions cancel each other so as to leave the pen at the point of beginning, unless  $\alpha = x\beta$ , i.e., unless  $\alpha$  is parallel to  $\beta$ .

32. If 
$$l\alpha+m\beta=l_1\psi+m_1\beta,$$
 that is, 
$$(l-l_1)\alpha+(m-m_1)\beta=0,$$
 then, 
$$l=l_1, \quad m=m_1.$$

- 33. If  $l\alpha + m\beta + n\gamma = 0$ , and l, m, n are not zero, then  $\alpha$ ,  $\beta$ , and  $\gamma$  are coplanar, for  $l\alpha$  and  $m\beta$  determine a plane which contains the *ends* of  $n\gamma$ , and therefore  $n\gamma$  itself.
- **34.** If  $l\alpha + m\beta + n\gamma = 0$ , and l, m, n are not zero but l+m+n=0, then  $(l+m+n)\alpha = 0$ , and subtracting the first equation, we get



$$m(\alpha-\beta)+n(\alpha-\gamma)=0,$$

whence  $\alpha - \beta$  and  $\alpha - \gamma$  are parallel (§ 31). But  $\alpha - \gamma$  connects the ends of  $\alpha$  and  $\gamma$ , and

 $\alpha - \beta$  the ends of  $\alpha$  and  $\beta$ , hence if l+m+n=0,  $\alpha$ ,  $\beta$ , and  $\gamma$  terminate in the same line.

**35.** Conversely, if  $\alpha$ ,  $\beta$ , and  $\gamma$  terminate collinearly and  $l\alpha + m\beta + n\gamma = 0$ , then l + m + n = 0.

For by condition,  $\alpha-\beta=x(\alpha-\gamma)$ , or  $(1-x)\alpha-\beta+x\gamma=0$ , in which 1-x-1+x=0. Q.E.D. **36.** If  $a\alpha + b\beta + c\gamma + \ldots = d\delta$ , then evidently  $a+b+c+\ldots > d$ , or  $TA + TB + TC + \ldots > TD$ ,

or, Sum of the tensors > tensor of the sum,  $\Sigma T > T\Sigma$ ,

or the distance a man travels > his distance from home.

If the vectors are parallel,  $\Sigma T = T \Sigma$ .

**37.** The diagonals of a parallelogram mutually bisect each other.

$$\alpha = \delta + y\gamma = \gamma + x\delta.$$


$$\therefore \S 31,$$

$$\delta = x\delta, \quad \gamma = y\gamma,$$

$$Q.E.D.$$

 $\gamma$  and  $\delta$  being parts of the diagonals to the point of intersection, and  $y\gamma$  and  $x\delta$  the remaining portions respectively.

**38.** The lines joining the middle points of the opposite sides of any quadrilateral, whether plane or gauche, mutually bisect each other.



One bisector is  $\alpha = \frac{1}{2}\lambda + \mu + \frac{1}{2}\nu$ .

The other is  $\beta = \frac{\mu}{2} + \nu + \frac{\rho}{2}$ .

Find the vectors from any assumed point to the middle points of these bisectors and

compare the results.


Thus the vector from the beginning of  $\lambda$  to the middle point of  $\alpha$  is


$$\zeta_1 = \frac{\lambda}{2} + \frac{\alpha}{2} = \frac{\lambda}{2} + \frac{1}{2} \left( \frac{1}{2} \lambda + \mu + \frac{\nu}{2} \right) = \frac{3}{4} \lambda + \frac{1}{2} \mu + \frac{1}{4} \nu.$$

The vector from the same point to the middle point of βis

$$\begin{split} &\zeta_2 = \lambda + \frac{\mu}{2} + \frac{\beta}{2} = \lambda + \frac{\mu}{2} + \frac{1}{2} \left( \frac{\mu}{2} + \nu + \frac{\rho}{2} \right) \\ &= \lambda + \frac{3}{4} \mu + \frac{1}{2} \nu - \frac{1}{2} \frac{\lambda + \mu + \nu}{2} = \frac{3}{4} \lambda + \frac{1}{2} \mu + \frac{1}{4} \nu. \end{split}$$

- $\therefore$   $\zeta_1 = \zeta_2$ , and the middle points coincide.
- Q.E.D.
- 39. If the ends of two parallel vectors be connected by 'straight lines, the join (connecting line) of the middle points of the straight lines is half the sum or difference of





the tensors of the parallel vectors: i.e., the median line of a trapezoid is half the algebraic sum of the bases.

$$\delta = \frac{\beta}{2} \pm \alpha - \frac{\gamma}{2},$$

taken along  $\alpha$ ;

$$\delta = -\frac{\beta}{2} + m\alpha + \frac{\gamma}{2}$$
, taken along  $m\alpha$ .

Whence by addition,  $\delta = \frac{m\alpha \pm \alpha}{2}$ .

$$d=\frac{ma\pm a}{2}$$
.

Q.E.D.

#### CHAPTER IV

#### MULTIPLICATION OF UNIT VECTORS

40. Parallel vectors. Remembering that multiplication is the performing by the reader on the multiplicand of an operation which is symbolized by the multiplier, viz., the operation which produced the multiplier from unity, we must in the product ii \* ask what operation is the first i the symbol of. The answer is, of course, of one of two equal operations whose successive applications shall produce reversal. Now i can be reversed by the repetition of each of two methods. The one we have designated as operation A (§ 20). The other we have designated as operation B. Since the multiplier is exactly the same as the multiplicand in all its properties, we must if possible use exactly the same operation not only in kind, but also in detail, that produced the multiplier, that is, operation B (§ 20).

This amounts to the repetition upon the multiplicand i of the same operation which produced it from unity, and of course results in -1, see § 20. That is,


ii = -1.

Hence, Multiplication (performance of an operation symbolized by the multiplier) of one unit vector into another parallel to it produces reversion.

<sup>\*</sup> i = some directed  $\sqrt{-1}$  = some directed mean reversed scalar.

41. Perpendicular vectors. Since the multiplier is now perpendicular to the multiplicand, we must take it as the symbol of an operation to be performed on the multiplicand, the same in kind but as far removed in detail from that which would have been used had the multiplier been parallel to the multiplicand as perpendicularity is removed from parallelism.

This we must do in order to take into account the perpendicularity of direction as opposed to parallelism. The



operation of mean reversion which produced the multiplier was operation B ( $\S$  20). Therefore we must use operation A. Let i and j be operator and operand respectively. The only position into which j can be revolved such that the reversal

of the signs of the two factors will give the same result is k, one of perpendicularity to both factors. Thus

$$ij=k$$
 and  $-i\cdot -j=k$ ,

since -i bears exactly the same relation to -j that i does to k, and must therefore have the same effect. Any other position than k for the product of ij would not do this.


Hence, The multiplication (performance of an operation symbolized by the multiplier) of one unit vector into another perpendicular to it results in the turning of the multiplicand through a right angle in (to) a plane perpendicular to the operator.

42. Inclined vectors. Naturally the result will be a combination of those of §§ 40, 41, that is partly scalar and partly vector, or

$$\alpha\beta = -\cos\theta + \varepsilon\sin\theta,$$

where  $\alpha$  and  $\beta$  are the two unit vectors inclined at an angle  $\theta$ , and  $\varepsilon$  a unit vector perpendicular to  $\alpha$  and  $\beta$ , since this formula satisfies both the limiting cases (§§ 40, 41).

Hence, The multiplication (performance of an operation symbolized by the operator) of one unit vector into another inclined to it at an angle  $\theta$ , thus producing the mean reversed



state induced by the operator symbol, turns the operand through a right angle into a plane perpendicular to the multiplier, makes its length  $\sin \theta$  and adds a scalar,  $-\cos \theta$ .

43. We can get the same result as follows:

 $i\beta$ , the mean reversed state of  $\beta$ , must be as to direction some vector perpendicular to the plane of i and  $\beta$ , since  $-i \cdot -\beta$  must produce the same result as  $i\beta$ . Hence, tentatively,

$$i\beta = sk$$
,

where s is some scalar. Operating again with i to see if the second application produces reversal, we get

$$i \cdot i\beta = i \cdot sk = s \cdot ik = s \cdot -j$$
,

which is not reversal, but which would be, except as to

length, perhaps, by the addition of -ci, where c is some scalar. But this would require

$$i\beta = -c + sk$$
, since  $i \cdot i\beta = -ic + i \cdot sk$   
=  $-ic - sj$ .

Now if  $-ic-sj=-\beta$ , then  $c=\cos\theta$ ,  $s=\sin\theta$ , and we have as before  $i\beta=-\cos\theta+k\sin\theta$ .

## 44. Exercises in unit reference vectors.

$$ij=k$$
, but  $ji=-k$ ;  $jk=i$ , but  $kj=-i$ .

Hence the factors are not commutative.

$$i \cdot jk = i \cdot i = -1,$$
  $ij \cdot k = k \cdot k = -1.$ 

Hence  $i \cdot jk = ij \cdot k$ , or the factors are associative.

$$ki = j,$$
  $i \cdot jk = ii = i^2 = -1,$   $ji = -k,$   $j \cdot ki = jj = -1,$   $i \cdot -j = -k,$   $ijk = jki = kij = -1,$   $ijk = jki = kij = -1,$   $k \cdot ji = k \cdot -k = -k^2 = 1.$   $ij \cdot k = kk = k^2 = -1.$ 

## CHAPTER V

### QUATERNIONS

**45.** Having ascertained that the product of two vectors  $\alpha$ ,  $\beta$  is

$$\alpha\beta = -\cos\theta + \varepsilon\sin\theta,$$

we can, § 29, express  $\epsilon \sin \theta$  in terms of i, j, k, viz.:

$$\varepsilon \sin \theta = xi + yj + zk$$
, or  $\alpha \beta = -\cos \theta + xi + yj + zk$ ,

which, being composed of four terms, a scalar and three vectors, is called a quaternion, and will be symbolized by q.

A quaternion is evidently composed of a scalar plus a vector. Later (§ 61) we shall find that, conversely, a scalar plus a vector is a quaternion.


46. The plane of the factors (multiplier and multiplicand) of a quaternion is called the plane of the quaternion.

The plane of a vector is the plane perpendicular to it.

 $\varepsilon$  is called the **axis** of  $\alpha\beta$ . The most convenient method of defining it seems to be: The unit vector toward the north pole when the multiplicand is to the east of the multiplier, the equator being the plane of the quaternion; toward the south pole when the multiplicand is to the west.

47. Meaning of  $\frac{1}{a}$ . By the rule for division we must first


ascertain what must be done to  $\alpha$  to produce 1. Returning to our idoigraphic shell; to convert  $\alpha$  into +1, we must, repeating the operation which produced it, apply operation B (§ 20), and then direct it one-half its dimen-



sions towards its correlatively reversed position. Performing these operations on the

#### numerator

we get by application of B, and then by directing it,



whence

$$\frac{1}{\alpha} = -\alpha$$
.

This is verified by the fact that  $\alpha \cdot -\alpha = -\alpha^2 = -\cdot -1 = 1$ , hence  $\frac{1}{\alpha} = -\alpha$ , since  $\alpha \frac{1}{\alpha} = 1$ , being a functional \* operation followed by the inverse operation and therefore resulting in the original operand.

Hence, the reciprocal of a unit vector is the unit vector reversed.

48. Meaning of  $\dagger \frac{\mathbf{j}}{\mathbf{i}}$ ,  $\frac{\mathbf{a}}{\mathbf{\beta}}$ , etc. In a similar manner  $\frac{\mathbf{j}}{\mathbf{i}} = k$ ,

<sup>\*</sup> See Appendix.

 $<sup>+\</sup>frac{j}{i}$  means  $j \cdot \frac{1}{i}$  and not  $\frac{1}{i} \cdot j$ . Thus we can write  $\frac{\alpha}{\beta} \cdot \frac{\beta}{\gamma} = \alpha \frac{1}{\beta} \beta \frac{1}{\gamma} = \alpha \frac{1}{\gamma}$ 

<sup>=</sup> $\frac{\alpha}{r}$ . But we cannot write  $\frac{\beta}{r} \cdot \frac{\alpha}{\beta} = \frac{\alpha}{r}$ , for  $\beta \frac{1}{r} \alpha \frac{1}{\beta} = \beta(-r)\alpha(-\beta)$  does not allow the  $\beta$ 's to cancel each other, the vector factors not being commutative, § 44.

and  $\frac{\alpha}{\beta} = \cos \theta - \varepsilon \sin \theta$ , since this satisfies both the limiting cases  $\frac{i}{i} = 1$  and  $\frac{j}{i} = +k$ ,  $\theta$  being the angle from  $\alpha$  to  $\beta$ .

# Examples for Practice

$$\begin{split} &\frac{-k}{i}=j, \quad \frac{ik}{j}=-1. & \qquad \frac{j}{k}=-i, \qquad \frac{i}{j}\frac{j}{k}=\frac{i}{k}. \\ &\frac{-i}{j}=k, \quad -i\cdot -j=k \qquad \qquad \frac{1}{i}=-i, \\ &\frac{-j}{k}=i, \quad i^2j^2k^2=-(ijk)^2, \quad \frac{j}{k}\cdot\frac{i}{j}=-i\cdot -k=-\cdot j=\frac{k}{i}\neq\frac{i}{k} \\ &\frac{j}{-k}=i, \quad \frac{i}{j}j=i, \qquad \qquad \frac{-j}{k}\cdot\frac{k}{j}=i\cdot -i=i^2. \\ &\frac{k}{j}=i, \quad j\frac{i}{j}\neq i, \qquad \qquad ikj=kji=jik=-i^2=-j^2=1. \end{split}$$

**49.** Since 
$$a\alpha \frac{1}{a\alpha} = \alpha \cdot -\alpha = 1 = \frac{1}{a\alpha} a\alpha = -\alpha \cdot \alpha$$
,

therefore a vector is commutative with its reciprocal.

**50.** Since 
$$\frac{1}{\beta} = -\beta$$
, we can write

$$\frac{1}{\beta}\beta = -\beta \cdot \beta = 1.$$


Hence, § 42,

$$\frac{\alpha}{\beta} = \alpha \cdot \frac{1}{\beta} = \alpha(-\beta)$$

$$= -\cos(\pi + \theta) + \varepsilon \sin(\pi + \theta)$$

$$= \cos\theta - \varepsilon \sin\theta,$$

where, as before,  $\theta$  is the angle from  $\alpha$  to  $\beta$ .



Similarly.

$$\beta \alpha = -\cos(-\theta) + \varepsilon \sin(-\theta) = -\cos\theta - \varepsilon \sin\theta$$

$$\frac{1}{\beta}\alpha = -\cos(\pi - \theta) + \varepsilon\sin(\pi - \theta) = \cos\theta + \varepsilon\sin\theta.$$

Hence

$$\frac{1}{\beta}\alpha \neq \left(\frac{\alpha}{\beta} = \alpha \frac{1}{\beta}\right),$$

$$-\frac{1}{\alpha}\beta = -\cos(\pi + \theta) + \varepsilon \sin(\pi + \theta) = \cos\theta - \varepsilon \sin\theta$$

$$\frac{\beta}{\alpha} = -\cos(\pi - \theta) + \varepsilon \sin(\pi - \theta) = \cos\theta + \varepsilon \sin\theta.$$

**51.** Introducing the tensors of  $\alpha$  and  $\beta$ , and collecting the results, we have,

$$\alpha\beta = ab(-\cos\theta + \varepsilon\sin\theta),$$
  $\frac{\alpha}{\beta} = \frac{a}{b}(\cos\theta - \varepsilon\sin\theta),$ 

$$\beta \alpha = ab \ (-\cos \theta - \varepsilon \sin \theta), \qquad \frac{\beta}{\alpha} = \frac{b}{a} (\cos \theta + \varepsilon \sin \theta).$$

**52.** Distributivity of the vector multiplier. Let  $\alpha$ ,  $\beta$ ,  $\gamma$  be three unit vectors, making with each other the angles  $\phi$ ,  $\theta$ , 2a, as shown;  $\delta$  is not a unit vector.

 $\varepsilon$  is the axis of  $\alpha\beta$ ,  $\eta$  of  $\alpha\gamma$ ,  $\zeta$  of  $\alpha\delta$ ; coplanar, since the the planes of the three quaternions have the common edge  $\alpha$ .

$$\beta + \gamma = \delta$$
,  $|\delta| = 2 \cos a$ .

[ $|\delta|$  means length of  $\delta$ .  $\cos a = \frac{1}{2}$  diag. of the parallelogram on  $\beta$ ,  $\gamma$ .]

$$\alpha\beta = -\cos\theta + \varepsilon\sin\theta,$$
  $\alpha\gamma = -\cos\phi + \eta\sin\phi.$ 


$$\therefore \quad \alpha\beta + \alpha\gamma = -\cos\theta - \cos\phi + \varepsilon\sin\theta + \eta\sin\phi.$$

But by trigonometry, since the angle between the axes is the same as the angle between the planes of the quaternions, and since a=a' makes the sines of these angles proportional to the sines of the adjacent sides, that is,  $\sin \theta$ ,  $\sin \phi$ , therefore  $\varepsilon \sin \theta + \eta \sin \phi$  will lie along the  $\zeta$  axis and

(1) 
$$\alpha\beta + \alpha\gamma = -\cos\theta - \cos\phi + x\zeta,$$

where x is some unknown tensor.

(2) But  $\alpha(\beta+\gamma) = \alpha\delta = 2\cos\alpha(-\cos\psi + \zeta\sin\psi)$ , and we now have to show that this agrees with  $\alpha\beta + \alpha\gamma$ .



By trigonometry,

$$\cos\phi = \cos\psi\cos a + \sin\psi\sin a\cos \Phi$$

$$\cos \theta = \cos \phi \cos a - \sin \phi \sin a \cos \Phi$$

$$\therefore \cos \Phi = \frac{\cos \phi - \cos \psi \cos a}{\sin \psi \sin a} = \frac{\cos \psi \cos a - \cos \theta}{\sin \psi \sin a},$$

$$\therefore \frac{\cos \phi + \cos \theta}{2 \cos a} = \cos \phi.$$

$$(3) \qquad \therefore \quad -2\cos a\cos \phi = -\cos \phi - \cos \theta.$$

By trigonometry again,

$$x^2 = \sin^2 \theta + \sin^2 \phi + 2 \sin \theta \sin \phi \cos A$$
,

 $\cos 2a = \cos \phi \cos \theta + \sin \phi \sin \theta \cos A$ .

$$\therefore x^{2} = \sin^{2}\theta + \sin^{2}\phi + 2(\cos 2a - \cos\phi \cos\theta)$$

$$= \sin^{2}\theta + \sin^{2}\phi + 2(\cos^{2}a - \sin^{2}a - \cos\phi \cos\theta)$$

$$= 1 - \cos^{2}\theta + 1 - \cos^{2}\phi + 2\cos^{2}a - 2 + 2\cos^{2}a$$

$$- 2\cos\phi \cos\theta$$

$$= 4\cos^{2}a - \cos^{2}\theta - 2\cos\theta \cos\phi - \cos^{2}\phi$$

$$= 4\cos^{2}a - (\cos\phi + \cos\theta)^{2}.$$

But

$$\sin \psi = \sqrt{1 - \left(\frac{\cos \phi + \cos \theta}{2 \cos a}\right)^2} = \frac{\sqrt{4 \cos^2 a - (\cos \phi + \cos^2 \theta)^2}}{\cos a}.$$

$$(4) \qquad \therefore \quad x = 2 \cos a \sin \phi.$$

$$\therefore \quad \alpha\beta + \alpha\gamma = -2\cos a \cos \phi + 2\cos a \sin \phi \cdot \zeta \quad (1) \cdot (3) \cdot (4)$$

$$= 2\cos a(-\cos \phi + \sin \phi \cdot \zeta)$$

$$= \alpha(\beta + \gamma) \cdot \qquad Q.E.D. \quad (2)$$

Hence in vector multiplication, the multiplier is distributive over the operand.

**53.** If 
$$\alpha\beta = ab(-\cos\theta + \varepsilon\sin\theta) = q$$
.

ab is called the **tensor** of q and is symbolized by ab = Tq.

- $-ab\cos\theta$  " scalar part of q and is symbolized by Sq.
- $ab \sin \theta \cdot \varepsilon$  " vector part of q and is symbolized by Vq.
- $ab \sin \theta$  "tensor of the vector part and is symbolized by TVq.
- $-\cos\theta + \epsilon\sin\theta$  " unit part and is symbolized by Uq.
  - $\epsilon$  "unit vector of the vector part and is symbolized by UVq.
  - $-\cos\theta$  " scalar of the unit part and is symbolized by SVq.
  - $\sin \theta \cdot \varepsilon$  "vector part of the unit part and is symbolized by VUq.
    - $\sin \theta$  "tensor of the vector part of the unit part and is symbolized by TVUq.

#### EXERCISES

# 54. Show that

$$(TVq)^2 = -(Vq)^2, \qquad \qquad UVq = \frac{Vq}{TVq},$$
 
$$(Tq)^2 = (Sq)^2 - (Vq)^2, \qquad \qquad TVUq = \frac{TVq}{Tq}.$$

If q=w+xi+yj+zk, § 45, show that Sq=w.

$$Vq = xi + yj + zk, \qquad (1) \qquad Tq = \sqrt{w^2 + x^2 + y^2 + z^2},$$

$$TVq = \sqrt{x^2 + y^2 + z^2}, \qquad Uq = \frac{w + xi + yj + zk}{\sqrt{w^2 + x^2 + y^2 + z^2}}.$$

$$UVq = \frac{xi + yj + zk}{\sqrt{x^2 + y^2 + z^2}},$$

If q and r are quaternions

(2) 
$$T \cdot qr = TqTr$$
,  $U \cdot qr = UqUr$ ,  $U\frac{q}{r} = \frac{Uq}{Ur}$ .

If q and r degenerate to  $\alpha$ ,  $T \cdot \alpha^2 = -a^2$ .  $[\alpha^2 = -1]$ .

**55.** By § 54, Eqs. (1), (2),

$$Tqr = \sqrt{\overline{W^2 + X^2 + Y^2 + Z^2}},$$

where

$$Tq = \sqrt{w_1^2 + x_1^2 + y_1^2 + z_1^2}, \quad Tr = \sqrt{w_2^2 + x_2^2 + y_2^2 + z_2^2}.$$

$$\therefore W^2 + X^2 + Y^2 + Z^2$$

$$= (w_1^2 + x_1^2 + y_1^2 + z_1^2)(w_2^2 + x_2^2 + y_2^2 + z_2^2),$$

or (Euler's Theorem) the sum of four squares may be resolved into two factors, each of which is the sum of four squares.

56. Show that

$$S\alpha\beta = S\beta\alpha, \qquad \alpha\beta - \beta\alpha = 2V\alpha\beta,$$

$$V\alpha\beta = -V\beta\alpha, \qquad \alpha\beta \cdot \beta\alpha = (S\alpha\beta)^2 - (V\alpha\beta)^2$$

$$\alpha\beta + \beta\alpha = 2S\alpha\beta, \quad (1) \qquad = (T\alpha\beta)^2.$$

$$(2) \qquad Sx = x, \quad Sij = 0, \qquad Vij = k.$$


**57.** Example. 
$$\alpha + \beta = \gamma$$
.

$$(\alpha + \beta)^2 = \gamma^2 = \alpha^2 + \alpha\beta + \beta\alpha + \beta^2,$$

or § 56, eq. (1), 
$$-a^2 + 2S\alpha\beta - b^2 = -c^2$$
.

If this is a right triangle, at C, then § 56, eq. (2),  $S\alpha\beta = 0$  and  $a^2 + b^2 = c^2$ .


If not a right triangle, this becomes



$$c^2 = a^2 + b^2 - 2ab \cos c$$
.

(Law of Cosines. Trig.)

# 58. Geometric meaning of TVaβ and Saβ.



$$V\alpha\beta = ab \sin \theta \cdot \varepsilon$$
,

$$TV\alpha\beta = ab \sin \theta$$

=parallelogram on  $\alpha\beta$ .

$$S\alpha\beta = -ab\cos\theta$$

= - (one tensor · projection of the other upon it).

**59.** Meaning of Sa $\beta\gamma$ . Suppose  $\alpha$ ,  $\beta$ ,  $\gamma$  unit vectors,  $\theta$  = angle between  $\alpha$ ,  $\beta$ , and  $\phi$  = angle between  $\gamma$  and plane of  $\alpha\beta$ .


$$\alpha\beta = -\cos\theta + \varepsilon\sin\theta,$$

$$S\alpha\beta\gamma = S\cdot(-\cos\theta + \varepsilon\sin\theta)\gamma$$

$$= S\varepsilon\gamma\sin\theta.$$

But  $S \varepsilon \gamma = -\sin \phi$ ,

$$\therefore S\alpha\beta\gamma = -\sin\phi\sin\theta.$$



$$S\alpha\beta\gamma = -abc\sin\theta\sin\phi$$

= -volume of the parallelopiped on  $\alpha$ ,  $\beta$ ,  $\gamma$ , as edges.

**60.** If  $S\alpha\beta\gamma=0$ ,  $\alpha$ ,  $\beta$ ,  $\gamma$  are coplanar, and vice versa.

61. The sum of a scalar and a vector is a quaternion. Let

$$w =$$
some scalar;

 $A = a\alpha =$ some vector.

Then

$$w + a\alpha = \sqrt{w^2 + a^2} \left( \frac{w}{\sqrt{w^2 + a^2}} + \alpha \frac{a}{\sqrt{w^2 + a^2}} \right)$$

=a tensor  $(\cos \phi + \alpha \sin \phi)$ 

62. Idiographic proof. We can always construct in a plane perpendicular to the vector a right-angle triangle, one of whose legs shall equal the diameter of the sphere which is the idiograph of the scalar, and the other leg the tensor of the given vector. If we denote the hypothenuse of this triangle by ab, then the leg corresponding to the scalar length will be  $ab \cos \theta$ , and the other leg will be  $ab \sin \theta$ ,  $\theta$  being the angle adjacent to the first leg constructed.

On the sides including  $\theta$ , lay off respectively the distances a and b as vectors. Then the product of these two vectors will be a quaternion,

$$-ab\cos\theta + ab\sin\theta \cdot \epsilon$$
,

which will be the given sum, and at the same time the product of two vectors.

63. We are now able to distinguish between the different branches of mathematics of discrete magnitudes.

First, the mathematics of numbers having size only. This is Arithmetic, the algebra of tensors. The operations are four only, addition, subtraction, multiplication, and division. The operands are tensors.

Secondly, the mathematics of numbers having size and sense, using in addition to the previous operations, *reversion*. Its operands are *scalars*, generally symbolized by letters and numerals. This is **Algebra**, in its broad sense, including Calculus and allied subjects.

The usual line of demarcation between arithmetic and algebra, the use or non-use of literal characters, is unphilosophical and erroneous in its significance. If the letters represent tensors (e.g., number of girls, boys, children), g+b=c is arithmetic purely, and 4-5=-1 is pure algebra.

Thirdly, the mathematics of numbers having size and sense, and which adds to the previous operations that of mean reversion. This is the algebra of Complex Functions. Its operands are scalars, symbolized by strokes as well as by letters and numerals.

Fourthly, that branch of mathematics which deals with numbers having size, sense, and direction. Its operands are scalars and vectors and combinations of these. This is the subject of Quaternions.

#### CHAPTER VI

#### KINDS OF QUATERNIONS

**64.** The reciprocal of a quaternion, denoted by Rq, is the factor into which the quaternion must be multiplied in order to produce unity.

If 
$$q = a\alpha b\beta$$
,  
then  $a\alpha b\beta \frac{1}{b\beta} \cdot \frac{1}{a\alpha} = \alpha\beta \frac{1}{\beta} \cdot \frac{1^*}{\alpha} = \alpha \frac{1}{\alpha} = 1$ .

Hence the reciprocal of a quaternion is found by taking the product, in reverse order, of the reciprocals of the factors.

$$\begin{split} R\frac{a\alpha}{b\beta} = b\beta\frac{1}{a\alpha}. & \frac{1}{\alpha\beta} = \frac{1}{q} = \frac{1}{-\cos\theta + \epsilon\sin\theta} = -\cos\theta - \epsilon\sin\theta. \\ & \frac{1}{\alpha}\cdot\frac{1}{\beta} = -\cos\theta + \epsilon\sin\theta. & \therefore & \frac{1}{\alpha\beta} \neq \frac{1}{\alpha}\cdot\frac{1}{\beta}. \\ & \frac{1}{\beta}\cdot\frac{1}{\alpha} = \frac{1}{\alpha\beta} \neq \frac{1}{\alpha}\cdot\frac{1}{\beta}. \end{split}$$
 If  $\alpha\beta = q$ , 
$$Rq = \frac{1}{\alpha\beta} = q^{-1}.$$

<sup>\*</sup> The assumption made here that  $\beta \frac{1}{\beta} \cdot \alpha = \beta \cdot \frac{1}{\beta} \alpha$  is justified later, in § 90, since all vectors are (§ 97) quaternions.

$$R\alpha\beta = \frac{1}{ab}(-\cos\theta - \varepsilon\sin\theta), \quad \frac{\alpha}{\beta} \cdot \frac{\beta}{\alpha} = 1, \quad \alpha\beta\frac{1}{\beta\alpha} \neq 1.$$

$$\frac{\alpha}{\beta} \cdot \frac{1}{\alpha} = \frac{\alpha}{\beta} \cdot \frac{1}{\alpha\frac{1}{\beta}} \neq \frac{\alpha}{\beta} \cdot \frac{1}{\alpha}\beta, \quad \alpha\beta \cdot \frac{1}{\alpha\beta} = q\frac{1}{q} = 1 = \frac{1}{\alpha\beta} \cdot \alpha\beta = \frac{1}{q} \cdot q.$$

Hence, a quaternion and its reciprocal are commutative:  $qq^{-1}=1=q^{-1}q$ .

65. Opposite quaternions are those in which one factor is reversed.

$$\alpha\beta = ab(-\cos\theta + \epsilon \sin\theta)$$

$$= ab(\cos(\pi - \theta) + \epsilon \sin(\pi - \theta)),$$

$$\alpha(-\beta) = ab(\cos\theta - \epsilon \sin\theta).$$
Hence if  $\alpha\beta = q$ ,  $\alpha(-\beta) = -q$ 
and  $q + (-q) = 0$ ,

or the sum of two opposite quaternions is zero.

$$\frac{-q}{q} = -1$$
,

or the quotient of two opposite quaternions is -1.

$$Tq = T(-q) = ab.$$

Opposite quaternions have a common plane, equal tensors, supplementary angles and opposite axes.

66. The Conjugate of a quaternion, denoted by Kq, has one of its factors turned an equal angle across the other factor, the order of the factors not being changed: it reverses the angle of the original quater-

Mon.  

$$Kq = K\alpha\beta = \alpha\beta'$$
  
 $= ab(-\cos(-\theta) + \varepsilon\sin(-\theta))$   
 $= ab(-\cos\theta - \varepsilon\sin\theta)$ .

$$Kq=Sq-Vq.$$
  $Kx=x.$   $q+Kq=2Sq.$   $\alpha\beta=K\beta\alpha.$   $q-Kq=2Vq.$  § 55.  $Kq=w-xi-yj-kz.$ 

Conjugate quaternions have a common plane, equal angles between the factors, equal angles (see § 76), equal tensors, and opposite axes. (§ 46.)

67. Since 
$$TKq = Tq$$
 and  $UKq = \frac{1}{Uq}$ .  

$$\therefore Kq = TKq \cdot UKq = \frac{Tq}{Uq}.$$

Whence by multiplication and division,

$$qKq = (Tq)^2,$$
  $\frac{q}{Kq} = (Uq)^2.$   $Kq = K\alpha\beta = Tq\frac{1}{Uq} = ab(-\cos\theta - \epsilon\sin\theta) = \beta\alpha.$   $\therefore K\alpha\beta = \beta\alpha \quad \text{and} \quad K\frac{\alpha}{\beta} = \frac{1}{\beta}\alpha.$ 

Evidently,  $Kxq = xab(-\cos \theta - \epsilon \sin \theta) = xKq$ . Making x = -1, we have

$$K(-q) = -Kq.$$

$$K\frac{1}{\alpha} = K\frac{\beta}{\alpha} = T\frac{\beta}{\alpha} \frac{1}{U\frac{\beta}{\alpha}} = T\frac{\beta}{\alpha} \cdot U\frac{\alpha}{\beta};$$

$$\frac{1}{K\frac{\alpha}{\beta}} = \frac{1}{T\frac{\alpha}{\beta}} \frac{1}{U\frac{\alpha}{\beta}} = T\frac{\beta}{\alpha}U\frac{\alpha}{\beta}; \qquad \therefore K\frac{1}{q} = \frac{1}{Kq}.$$

Confirm this by diagram, using  $\alpha\beta$  instead of  $\frac{\alpha}{\beta}$ .

**69.** Collecting the q,  $q^{-1}$ , -q, and Kq into a table, we have:

			T	$\theta_1$	U
$\frac{\alpha}{\beta}$	q	$\frac{a}{b}(-\cos\phi+\epsilon\sin\phi)$	Tq	φ	Uq
$\frac{\beta}{\alpha}$	q-'	$\frac{b}{a}(-\cos\phi-\epsilon\sin\phi),$	$rac{1}{Tq}$	$-\phi$	$\frac{1}{Uq}$
$\frac{-\alpha}{\beta}$	-q	$\frac{a}{b}(\cos\phi - \varepsilon\sin\phi)$	Tq	π+•φ	$\frac{1}{Uq}$
$\frac{1}{\beta}\alpha$	Kq	$\frac{a}{b}(-\cos\phi - \epsilon\sin\phi)$	$Tq^{ullet}$	-φ	$rac{1}{Uq}$

 $\theta_1$  indicates the angle from the multiplier to the multiplicand, and not the angle between  $\alpha$  and  $\beta$ .

From this table it is evident that if  $\phi = 0$ , KS = S, i.e., The conjugate of a scalar is the scalar itself.

If 
$$\phi = \frac{\pi}{2}$$
,  $KV = -V$ , The conjugate of a vector is its opposite.

70. 
$$RRq = \left(\frac{1}{q}\right)^{-1} = q$$
, The reciprocal of the reciprocal

$$-(-q) = q$$
, The opposite of the opposite

$$KKq = q$$
, The conjugate of the conjugate of the conjugate

71. Equality of quaternions. If  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$  are four coplanar unit vectors with the same angle  $\theta$  between  $\alpha$  and  $\beta$ , and  $\gamma$  and  $\delta$ , then

$$\alpha\beta = -\cos\theta + \varepsilon\sin\theta, \qquad \gamma\delta = -\cos\theta + \varepsilon\sin\theta,$$
 or 
$$\alpha\beta = \gamma\delta.$$
 Similarly, 
$$\frac{\alpha}{\beta} = \frac{\gamma}{\delta} = \cos\theta - \varepsilon\sin\theta.$$


1

Hence, Revolving the factors of a quaternion in the plane of the quaternion does not alter the quaternion.

Quaternions having the same S and V parts must be the same. Diplanar quaternions, that is, quaternions not having their factors in the same plane, cannot have the same V part, and therefore cannot be equal. To be equal, two quaternions must be coplanar.

72. Two quaternions can always be transformed so as to have the same vector for the numerator of one and the denominator of the other, or what is the same thing, the multiplicand of one the multiplier of the other.

For moving the quaternions in their planes until the denominator of one and the numerator of the other lie



along the line of intersection of the planes, a vector along this line can be taken as the numerator of one quaternion quotient and the denominator of the other. Thus

$$\frac{\alpha}{\beta} = \frac{\mu}{\nu}. \qquad \qquad \frac{\gamma}{\delta} = \frac{x\nu}{\rho} = \frac{\nu}{\frac{\rho}{x}}. \qquad Q.E.D$$

#### CHAPTER VII

### THE QUATERNION AS A MULTIPLIER

73. Into the reciprocal of its multiplicand factor.

$$\frac{\alpha}{\beta}\beta = q\beta = \alpha$$
.

Result: The quaternion multiplier turns the multiplicand through the angle  $-\theta$ , in the plane

74. Into its mulitplier factor.

of the quaternion.

$$q\alpha = \frac{\alpha}{\beta}\alpha = \alpha \cos \theta - \epsilon \alpha \sin \theta$$
$$= \alpha \frac{1}{\beta}\alpha = \alpha \cos \theta + \alpha \epsilon \sin \theta.$$

[§ 51.


a cos

Result: Same as in § 73.

**75.** Into any coplanar factor, γ.


$$\frac{\alpha}{\beta}\gamma = \frac{\alpha}{\beta}(-x\alpha + y\beta) = -x\alpha \cdot \frac{1}{\beta}\alpha + y\frac{\alpha}{\beta}\beta$$

$$= -x\alpha(\cos\theta + \varepsilon\sin\theta) + y\alpha = -x\alpha\cos\theta - x\sin\theta \cdot \alpha\varepsilon + y\alpha.$$



Result: Same as in § 73.

76. Into any quaternion. Let the quaternions be reduced, § 72, to a common numerator and denominator respectively, viz.:



$$\frac{\alpha}{\beta} \cdot \frac{\beta}{\gamma} = \frac{\alpha}{\gamma}$$
.

Denote the arcs which determine the planes of the quaternions by c, a, b respectively, as shown in the figure. The effect of the quaternion whose axis is  $ax \cdot c$  operating upon the quaternion whose axis is  $ax \cdot a$  is to give a new quaternion whose axis is  $ax \cdot b$ .

This can be analyzed as follows: Suppose we move the arc a horizontally along c, being careful not to change its inclination to c, until it takes the position a'. The  $ax \cdot a$  will move horizontally around  $ax \cdot c$  (conical revolution) until it takes the position  $ax \cdot a'$ . Then as we turn a' down to coincidence with b around a as an axis,  $ax \cdot a'$  will move up the arc passing through  $ax \cdot c$  (the equator of a) until it takes the position  $ax \cdot b$ .

That is, the multiplication of  $\frac{\alpha}{\beta}$  into  $\frac{\beta}{\gamma}$  turns the axis of  $\frac{\beta}{\gamma}$  horizontally, i.e., parallel to the plane of  $\frac{\alpha}{\beta}$  (conical revolution) through an angle equal to that between  $\alpha$  and  $\beta$ , and raises or lowers it through an angle which depends entirely upon  $\frac{\beta}{\gamma}$ . Whatever the position of  $\gamma$  with reference to  $\beta$ , the horizontal revolution is always the same, dependent upon  $\alpha$  and  $\beta$ .

Hence, the effect of  $\frac{\alpha}{\beta}$  as a multiplier is generally to revolve the axis of the multiplicand conically through a definite angle around the axis of the multiplier. This angle  $(-\theta)$  here, the angle through which the axis of the operand is moved, is called the **angle of the quaternion**, and is designated by  $\angle q$ , or in these notes by D.

77. Since the effect of a quaternion, q, as a multiplier is to revolve the multiplicand (axis) through the angle  $\angle q$  around the  $ax \cdot q$ , the arc of a great circle which measures  $\angle q$  on the equator of  $ax \cdot q$ , can be taken as an indirect measure of Uq.

Just as a quaternion can be revolved in its plane without alteration of value, so the arc representing it can be moved in its *great circle* without alteration of value.

Arcs in different great circles cannot be equal. If they are semicircles, any semicircle will have the same effect.


78. Representing the quaternions  $\frac{\alpha}{\beta}$ ,  $\frac{\beta}{\gamma}$ ,  $\frac{\alpha}{\gamma}$ , by their arcs (arc strokes) c, a, b respectively, we have, considering the arcs as arc strokes on the sphere surface similar to the plane strokes of § 12,

$$\frac{\alpha}{\beta} \cdot \frac{\beta}{\gamma} = \frac{\alpha}{\gamma}$$
 or  $a+c=b$ ,


and we have reduced the multiplication of quaternions to the addition of arc strokes.

In the same way

$$\frac{\beta}{\alpha} \cdot \frac{\alpha}{r} = \frac{\beta}{r}, \quad b + (-c) = a.$$



- 79. Notice carefully that the arc strokes corresponding to the quaternion factors read from the left, are added up from the right.
- 80. Evidently from the figure qq' is quite a different arc from q'q, and hence, The addition of arc strokes is not commutative, unless they are coplanar; and correspondingly, the multiplication of quaternions is not commutative, unless the quaternions are coplanar.



Show that

Rule for multiplication by arc strokes. To the arc stroke of the multiplicand add the arc stroke of the multiplier.

Rule for division by arc strokes. Reverse the arc stroke of the divisor and add the arc stroke of the dividend.

$$qKq = Kq \cdot q. \qquad q(-q) = -q \cdot q.$$
 
$$q\frac{1}{q} = \frac{1}{q}q = qq^{-1} = q^{-1}q.$$

81. From this we see that the product of a quaternion, is a quaternion, the sum of two arc strokes being an arc stroke.

Take two quaternions in the form of a scalar plus a vector, and show that their product is a scalar plus a vector, and therefore, § 61, a quaternion.

**82.** By multiplying the various forms of a quaternion,  $\alpha\beta$ ,  $\beta\alpha$ ,  $\frac{\alpha}{\beta}$ ,  $\frac{\beta}{\alpha}$ , etc., into some vector, whatever will give the result most readily, we find the following angles for the quaternions,  $\theta$  being the angle from  $\alpha$  to  $\beta$ ,  $\theta_1$  the angle from the multiplier to the multiplicand, and D the angle of the quaternion.


		$\theta_1$
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	θ	$\pi - \theta$
	<b>-</b> θ	$\pi + \theta$
	$-\theta$	$-\pi+\theta$
$\frac{eta}{lpha}$ $ heta$ $ heta$		

Inspection of the table shows that: The angle, D, of a quaternion is the supplement of the angle between the factors measured from the multiplier to the multiplicand.

83. Representing the angle measured from  $\alpha$  to  $\beta$  by  $\theta$ , and the angle of the quaternion by D, the angle of  $\alpha\beta$  by  $\phi$ , the angle of  $\frac{\alpha}{\beta}$  by  $\phi$ , and abbreviating  $\cos D + \varepsilon \sin D$  by  $\cos D$ , we get the following table:

		l		
$\alpha \beta$	cεs ψ	$ces(\pi+\phi)$	$c \varepsilon s (\pi - \theta)$	Ces D
βα	$\cos (2\pi - \psi)$	$ces(\pi-\phi)$	$c \in s(\pi + \theta)$	ces D
$\frac{\alpha}{\beta}$	$c \in s(\pi + \psi)$	cεs φ	cεs (-θ)	ces D
$\frac{\beta}{\alpha}$	$c \in (\pi - \psi)$	$\cos (2\pi - \phi)$	cεs θ	ces D
$\frac{1}{\beta}\alpha$	$c \in (\pi - \psi)$	<b>c</b> εs ( – φ)	cεs θ	ces D
$\frac{1}{\alpha}\beta$	$c \in (\pi + \psi)$	cεs φ	cεs (-θ)	ces D

84. Since a vector is the special case of a quaternion, one whose angle is  $\frac{\pi}{2}$ , we can represent a vector by an arc stroke



of  $\frac{\pi}{2}$  on its equator. As an example

take the case  $\frac{\alpha}{\beta}\gamma$ .

$$\operatorname{arc} \frac{\alpha}{\beta} = c$$
,  $\operatorname{arc} \gamma = a$ ,

$$\frac{\alpha}{\beta}\gamma = a + c = b = \text{arc } \delta,$$

and  $\gamma$  has been revolved through  $\angle q$ .

85. Collecting the results of these articles we have the table on the following page, where

$$\theta_1 = \pi - D, \qquad D = \angle q, \qquad \psi = \angle \alpha \beta, \qquad \phi = \angle \frac{\alpha}{\beta} = \angle \frac{1}{\alpha} \beta,$$

$$\theta = \angle \frac{\beta}{\alpha} = \angle \frac{1}{\beta} \alpha$$
.  $\theta_1$  = angle from first factor to the second.

 $\theta$  = angle from  $\alpha$  to  $\beta$ .

86. Since quaternions are multiplied by adding their arc strokes, the square,  $q^2$ , of a quaternion will have for its representative arc stroke twice the stroke of q. Similarly, one-half the arc stroke will be representative of  $q^{\frac{1}{2}}$ . But since there are two arcs D and  $-(2\pi-D)$ ,  $q^{\frac{1}{2}}$  will have two representative strokes,  $\frac{1}{2}D$  and  $-\pi + \frac{D}{2}$ , either of which if doubled will give the stroke of q. Hence, Rule for extraction of square roots of unit quaternions: Halve the angles of the quaternion.

$b_A$	$ab \sin \theta \cdot \epsilon$	$-ab\sin\theta\cdot \epsilon$	$-\frac{a}{b}\sin\theta \cdot \epsilon$	$\frac{b}{a}$ sin $\theta \cdot \epsilon$	$-\frac{b}{a}\sin\theta\cdot\epsilon$	$\frac{a}{b}\sin\theta \cdot \epsilon$	$ab \cos \theta - ab \sin \theta \cdot \epsilon$	
bs:	$-ab\cos\theta$	$-ab\cos\theta$	$\frac{a}{b}\cos\theta$	$\frac{b}{a}\cos\theta$	$\frac{a}{b}\cos\theta$	$\frac{b}{a}\cos\theta$	ab cos θ	
Rq	$\frac{1}{\beta \alpha}$	$\frac{1}{\alpha \beta}$	β	ष्ठ	$\frac{1}{\beta}\alpha$	$\frac{1}{\alpha}\beta$	$\frac{-1}{\beta \alpha}$	
	φ+μ	φ-μ	\$	φ 	\$	ф 	ъ.	
<b>b</b> 7	æ	-¢-	n+ 4	я— н ф— н	n + φ	$\pi - \phi$	n+4	
	$\mu - \mu$	$\theta + \omega$	θ-	0	θ-	θ	θ-	κ Q
Tq	ab	ap	ماه	عام	عام	ماه	ę	П
D of ces D	$\mu - \mu$	$\mu + \theta$	θ-	θ	θ-	θ	θ-	K   63
$U_{oldsymbol{q}}$	$-\cos \theta + \sin \theta$	$-\cos \theta - \epsilon \sin \theta$	$\cos(-\theta) + \epsilon \sin(-\theta)$	$\cos \theta + \epsilon \sin \theta$	$\cos \theta - \epsilon \sin \theta$	$\cos \theta + \epsilon \sin \theta$	$\cos \theta - \epsilon \sin \theta$	w
0,	θ	θ-	$\mu + \theta$	$\mu - \mu$	$\theta + \mu$	$\mu - \theta$	$u+\theta$	# C
ō	αġ	βα	ø ø.	B	$\frac{1}{\alpha^{\prime}\beta}$	$\frac{1}{\beta}$	- αβ	•

87. Since the angle of the quaternion is the supplement of the angles measured from the multiplier to the multiplicand, that is,  $\pi - D = \theta_1$ , a better working rule would be:

For square root. Revert the multiplicand and halve the angles between this and the multiplier, for the factors of  $q^{\frac{1}{2}}$ .

Similarly, a working rule for squaring quaternions: Double the angle between the factors and revert the multiplicand in its new position for the factors of  $q^2$ .

In  $\delta^2 = \beta \gamma$ , or  $\frac{\delta}{\beta} = \sqrt{\frac{\gamma}{\beta}}$ ,  $\delta$  bisects the angle between  $\beta$  and  $\gamma$ .

88. Using the arcs of quaternions, show that

- (1)  $\operatorname{arc} q^2 = \operatorname{arc} q + \operatorname{arc} q = 2 \operatorname{arc} q \pm 2n\pi.$
- (2)  $\operatorname{arc} q^{\frac{1}{2}} = \frac{1}{2} (\operatorname{arc} q \pm 2n\pi).$
- (3)  $\operatorname{arc} q^{n} = \operatorname{arc} q + \operatorname{arc} q + \ldots = n \operatorname{arc} q \pm 2n\pi.$

(4) 
$$\operatorname{arc} q^{\frac{1}{n}} = \frac{1}{n} (\operatorname{arc} q \pm 2n\pi).$$
 (5)  $(pq)^2 \neq p^2 q^2.$ 

89. When the quaternion is a scalar, its arc is some multiple of  $\pi$ . All coinitial arcs of the same length,  $\pi$ , for instance, will represent will represent the same scalar, -1, however the arcs are situated; that is, whatever the direction of their axes. In halving these arcs to extract square roots we will get an infinite number of answers, one for each arc, each corresponding to its own axis. Or in other words, there are an infinite number of vectors the square of which is a scalar.

Similar reasoning shows that there are an infinite number of quaternion *n*th roots of a scalar. On this account the roots of a scalar are limited to scalars.

From Eq. (2) of § 88 we get  $(pq)^{\frac{1}{2}} = \cos \frac{D}{2} + \varepsilon \sin \frac{D}{2}$ ; repetition of which gives  $(pq)^{\circ} = \cos 0 = 1$ , thus illustrating for the quaternion number the familiar fact of algebra.

#### CHAPTER VIII

### PRODUCTS OF QUATERNIONS

**90.** Having three quaternions, let the first two be reduced, § 72, to a common denominator and numerator respectively, viz.,  $q = \frac{\alpha}{\beta}$ ,  $r = \frac{\beta}{\gamma}$ ; and the latter two also, viz.,  $r = \frac{\mu}{\nu}$ ,  $s = \frac{\nu}{\rho}$ , where, evidently, since they represent the same quaternion,

$$\frac{\beta}{\gamma} = \frac{\mu}{\nu}$$
.

The product of the three quaternions is

$$\frac{\alpha}{\beta} \left( \frac{\beta}{\gamma} = \frac{\mu}{\nu} \right) \frac{\nu}{\rho} = qrs.$$

Associating the first two factors, we have,

$$qr \cdot s = \frac{\alpha}{\beta} \frac{\beta}{\gamma} \cdot \frac{\nu}{\rho} = \frac{\alpha}{\gamma} \cdot \frac{\nu}{\rho} = \alpha \cdot \frac{1}{\gamma} \nu \cdot \frac{1}{\rho}.$$

Associating the second two, we have

$$q \cdot rs = \frac{\alpha}{\beta} \left( \frac{\mu}{\nu} \cdot \frac{\nu}{\rho} \right) = \frac{\alpha}{\beta} \cdot \frac{\mu}{\rho} = \alpha \cdot \frac{1}{\beta} \mu \cdot \frac{1}{\rho}.$$

But if 
$$\frac{\beta}{\gamma} = \frac{\mu}{\nu}$$
 then  $\frac{\gamma}{\beta} = \frac{\nu}{\mu}$ .

Multiplying this by  $\frac{1}{r}$  and into  $\mu$ , we have

$$\frac{1}{\beta}\mu = \frac{1}{\gamma}\nu.$$

Whence


$$\frac{\alpha}{\gamma} \cdot \frac{\nu}{\rho} = \frac{\alpha}{\beta} \cdot \frac{\mu}{\rho}$$
 or  $qr \cdot s = q \cdot rs$ ,


or quaternions are associative in multiplication.

91. Since, § 67,

$$Kq = \frac{Tq}{Uq} = Tq(-\cos\theta - \epsilon\sin\theta) = Tq(-\cos(-\theta) + \epsilon\sin(-\theta)),$$

the arc representing Kq will be the reverse of that representing q, or  $xq = -\operatorname{arc} q$ .





Inspection of the figures shows that

$$-\operatorname{arc} qr = +\operatorname{arc} KrKq.$$

Hence

$$K(qr) = KrKq,$$
 or

The conjugate of the product of two quaternions is the product of their conjugates in a reverse order.

$$K(\alpha\beta) = -\beta \cdot -\alpha = \beta\alpha$$
. Conf. § 67.

**92.** Coplanar quaternions. Using arc strokes for q and r show that

$$qr = rq$$
. Conf. § 80.

Therefore, since a quaternion and its reciprocal, opposite, conjugate, and any power are coplanar,

$$qKq = Kq \cdot q$$
.  $q^{-1} \cdot -q = -q \cdot q^{-1}$ .  $Kq \cdot q^n = q^nKq$ .

**93.** Using q = Sq + Vq, r = Sr + Vr, show that qr is a quaternion, the sum of a scalar and a vector.

Show that

$$\begin{split} Sqr = &SqSr + S \cdot VqVr. & V \cdot VqVr = -V \cdot VrVq. \\ Vqr = &SqVr + SrVq + V \cdot VqVr. & qr \neq rq. & \text{unless coplanar.} \\ Srq = &SrSq + S \cdot VrVq. & T \cdot qr = TqTr. \\ Vrq = &SrSq + SqSr + V \cdot VrVq. & V \cdot qr = VqVr. \\ Srq = &Sqr. & S \cdot qr \neq SqSr, \end{split}$$

unless  $S \cdot VqVr = 0$ , i.e., unless the planes of the two quaternions are perpendicular.

$$Vqr \neq Vrq$$
,

unless  $V \cdot VqVr = 0$ , i.e., unless the quaternions are coplanar. Since

$$(p+q)(r+s) = pr + qr + ps + qs = pr + ps + qr + qs = etc.,$$

therefore the distributive and associative law applies to quaternions, verified by resolving the quaternions into their vector factors and applying the results of § 52.

### CHAPTER IX

#### **VERSORS**

94.  $U\alpha\beta = -\cos\theta + \epsilon\sin\theta = \cos D + \epsilon\sin D$ , turns any quaternion to which it is applied as a multiplier, horizontally through the angle D.

If  $D = \frac{\pi}{2}$ , then  $U\alpha\beta = \epsilon^1$  turns the multiplicand quaternion

through the angle  $\frac{\pi}{2}$ . Applied again it turns it through another right angle, or in all through the angle  $\pi$ . That is  $\epsilon$  turns the multiplicand through one rt.  $\angle$ ,  $\epsilon^2$  turns it through two rt.  $\angle$ 's, and so on.\* Hence, by the law of mathematical continuity,  $\epsilon^{\frac{1}{2}}$  should be the symbol of turning

through  $\frac{1}{2}$  rt.  $\angle$ , etc. If  $\phi$  is expressed in radians,  $e^{\frac{2}{\pi}\phi}$  turns through the angle  $\phi$ .

If  $\epsilon$  is perpendicular to its operand, we recognize the familiar equations,

$$ij = k$$
,  $i^2j = -j$ ,  $i^3j = -k$ , etc.

According to this notation then,

$$U\alpha\beta = \cos D + \varepsilon \sin D = \varepsilon^{\frac{2}{\pi}D}.$$

<sup>\*</sup>Remember that  $\epsilon^2$  does not mean here  $\epsilon$  multiplied by itself, but  $\epsilon$  applied twice to some operand. See Appendix.

95. Since  $\epsilon$  and  $-\epsilon$  are opposed,  $\epsilon^{-\frac{2}{\pi}D}$ , which turns clockwise from the upper side of the plane, must have the same effect as  $(-\epsilon)^{\frac{2}{\pi}D}$ , which turns counterclockwise from the lower side of the plane. Hence

$$\varepsilon^{-\frac{2}{\pi}D} = (-\varepsilon)^{\frac{2}{\pi}D}$$
, or  $e^{-\beta} = (-\varepsilon)^{\beta}$ .

If  $D = \frac{\pi}{2}$ , then  $e^{-1} = -e$ , or to put it in the more familiar form,  $i^{-1} = -i$ .

96. Again, § 85,

$$U^{\beta}_{\alpha} = \cos \theta + \varepsilon \sin \theta = \varepsilon^{\frac{2}{\pi}\theta},$$

and

$$U^{\alpha}_{\overline{\beta}} = \cos(-\theta) + \epsilon \sin(-\theta) = \epsilon^{-\frac{2}{\pi}\theta};$$

therefore,  $\varepsilon^{-\theta} = \frac{1}{e^{\theta}}$ .

From these 
$$e^{-\theta} = \cos(-\theta) + \epsilon \sin(-\theta) = (-\epsilon)^{\theta}$$
,  
 $-(\epsilon^{\theta}) = -\cos\theta - \epsilon \sin\theta$ .

Hence in general,

$$(-\varepsilon)^{\theta} = \varepsilon^{-\theta} \neq -(\varepsilon^{\theta})$$

unless  $\theta = 1$ , in which case, as before,  $\epsilon^{-1} = -\epsilon$ .

97. The unit portion of a quaternion (considered as the symbol of an operation to be performed by the reader) results in the turning of the operand, and is therefore aptly termed a versor.

It must not be forgotten that all vectors (considered as symbols of an operation to be performed by the reader) are versors, as indeed all vectors are special cases of quaternions. The evanescence of the scalar part does not affect its versorial character.

- i, j, and k are called quadrantal versors.
- 98. The method of expressing the unit part of a quaternion as the power of a vector gives a convenient method of indicating multiplication and division of coplanar quaternions.

$$Uq = \cos \phi + \varepsilon \sin \phi = \varepsilon^{\frac{2}{\pi}\phi}. \qquad Uq' = \cos \psi + \varepsilon \sin \psi = \varepsilon^{\frac{2}{\pi}\phi}.$$

$$\therefore UqUq' = (\cos \phi + \varepsilon \sin \phi)(\cos \psi + \varepsilon \sin \psi) = \varepsilon^{\frac{2}{\pi}\phi}\varepsilon^{\frac{2}{\pi}\phi}.$$

$$= \cos (\phi + \psi) + \varepsilon \sin (\phi + \psi) \qquad = \varepsilon^{\frac{2}{\pi}}(\phi + \psi).$$
Similarly,
$$\varepsilon^{\phi} \cdot \varepsilon^{-\phi} = \varepsilon^{\phi}\varepsilon^{\phi} = \varepsilon^{\phi - \phi},$$
and
$$(\varepsilon^{\phi})^{m} = \varepsilon^{m\phi}.$$

Hence, The algebraic law of indices holds good for versors. Hence, The angle of the product of two coplanar quaternions is the sum of the angles of the two factors.

99. According to this notation,

$$\alpha\beta = 4(-\cos 30^{\circ} + \epsilon \sin 30^{\circ})$$

$$= 4(\cos 150^{\circ} + \epsilon \sin 150^{\circ}) = 4\epsilon^{\frac{5}{3}}.$$

$$\alpha\beta = 2(\cos 60^{\circ} - \epsilon \sin 60^{\circ})$$

$$= 2(\cos (-60^{\circ}) + \epsilon \sin (-60^{\circ})) = 2\epsilon^{-\frac{5}{3}}.$$

Show that 
$$q^2 = \{Tq(\cos\phi + \epsilon\sin\phi)\}^2 = Tq)^2 \epsilon^{\frac{4\phi}{\pi}}$$
  
=  $(Tq)^2(\cos 2\phi + \epsilon\sin 2\phi)$ .  
 $q^n = (Tq)^n(\cos n\phi + \epsilon\sin n\phi)$ .

Hence any quaternion can be written, where  $\rho$  is some vector,

$$\rho^t = \rho^{\frac{2\phi}{\pi}}$$
.

**100.** In the diagram to § 57,

$$\frac{2A}{\varepsilon^{\frac{2}{\pi}}}\gamma = \beta. \qquad \varepsilon^{\frac{2B}{\pi}}\alpha = \gamma. \qquad \varepsilon^{\frac{2C}{\pi}}\beta = -\alpha.$$

$$\therefore \varepsilon^{\frac{2(A+B+C)}{\pi}} = -1 \quad \text{or} \quad A+B+C=\pi.$$

101. What will convert rq into qr? Assume for the moment that the tensors are unity. Let the strokes of q and r be as shown. Then the strokes of qr and rq will take the positions shown.

From the table of § 83, we find

$$q=rac{lpha}{eta}, \qquad r=rac{\gamma}{eta}, \qquad qr=rac{lpha}{\gamma}, \qquad rq=rac{\gamma'}{lpha'},$$

and the problem becomes: To con-

vert 
$$\frac{\gamma'}{\alpha'}$$
 into  $\frac{\alpha}{\gamma}$ . Now

$$\frac{\alpha}{\gamma} = \frac{\alpha}{\beta} \cdot \frac{\beta}{\gamma'}, \qquad = \frac{\alpha}{\beta} \cdot \frac{\gamma'}{\beta}, \qquad \S 71.$$


$$= \frac{\alpha}{\beta} \cdot \frac{\gamma'}{\alpha'} \cdot \frac{\alpha'}{\beta}, \qquad = \frac{\alpha}{\beta} \cdot \frac{\gamma'}{\alpha'} \cdot \frac{\beta}{\alpha}. \qquad \S 71.$$

Or substituting for these their equivalents, § 69,

$$qr = q(rq)q^{-1}$$
.

102. Meaning of  $q(r)q^{-1}$ . Indicate q and r by their arc strokes, as shown in the diagram. The other strokes will have the values indicated.

The two triangles have two sides and the included angle



respectively equal, and, as shown in the figure, by the angle  $\theta$ ,  $qrq^{-1}$  is r revolved in the plane of q through  $2 \angle q$ . This amounts to revolving the axis of r conically around the axis of q through  $2 \angle q$ .

Hence  $qrq^{-1}$  differs from r only in being rotated through a certain angle. Hence

 $q()q^{-1}$  may be aptly called a rotator, since it rotates any quaternion inserted in the parenthesis.

This is a special case of a more general function, called a nonion, which we shall meet with farther on.

103. If r is some multiple of q, say q'q, then

$$q(q'q)q^{-1} = qq'qq^{-1} = qq'.$$

Hence qq' is q'q rotated through the angle  $2 \angle q$  in the plane of q.

Compare this with the figure of § 80, and see how they agree.

Similarly,  $q^{-1}rq$  rotates r negatively through  $2 \angle q$ .

Since, 
$$Tq \cdot T \frac{1}{q} = 1$$
, therefore,  $T \cdot qrq^{-1} = Tr$ ,

so that in the discussion above only the unit or versor parts needed to be considered.

$$q\beta q^{-1} = \beta',$$

revolves the plane of  $\beta$ , or  $\beta$ , conically around the axis of q through  $2 \angle q$ .

105.

$$\alpha\beta\alpha^{-1}=\beta'$$
,

revolves  $\beta$  conically around  $\alpha$  through the angle  $2\pi$ , that is, turns  $\beta$  in the plane of  $\alpha\beta$  to a corresponding position on the other side of  $\alpha$ .


**106.** Exercises. If  $\alpha \gamma = q$ , show that  $\alpha q \alpha^{-1} = Kq$ . If  $\alpha, \beta, \gamma$  are coplanar, show that  $\alpha \cdot \beta \gamma \cdot \alpha^{-1} = \alpha^{-1} \beta \gamma \alpha = K \beta \gamma$ .

$$\begin{split} (qrq^{-1})^{x} = &qr^{x}q^{-1}. & (jkj^{-1})^{2} = &jk^{2}j^{-1}. \\ & rqBq^{-1}r^{-1} = &rq\cdot B\cdot (rq)^{-1}. \end{split}$$

#### CHAPTER X

#### INTERPRETATION OF VECTOR EQUATIONS

107.  $\alpha\beta\gamma = \alpha\beta^2\beta^{-1}\gamma = -b^2\frac{\alpha}{\beta}\gamma = \delta$ ,  $\alpha$ ,  $\beta$ ,  $\gamma$  coplanar, i.e.,  $S\alpha\beta\gamma = 0$ .  $U \cdot -\frac{\alpha}{\beta}$  acting on  $\beta$  turns it through an angle  $\theta$  into  $-\alpha$ . Acting on  $\gamma$  it turns it through the same angle into  $\delta$ . If  $\alpha$ ,  $\beta$ ,  $\gamma$  represent in direction the successive sides of a polygon (which is always possible) then  $U \cdot -\frac{\alpha}{\beta}$  must turn  $\gamma$  into a direction coinciding with the fourth side of



a polygon inscribed in a circle drawn through the intersection of  $\alpha$ ,  $\beta$ , and  $\gamma$ . The diagram shows this. Hence  $\delta$  is in direction the fourth side of the inscribed polygon of which  $\alpha$ ,  $\beta$ , and  $\gamma$  are the other three.

If the circle passes through three intersections, that circumscribes  $\alpha$   $\beta$ ,  $\gamma$  as a triangle, then  $\delta$  becomes tangent to this circle.

108. Locus of  $\xi$  in  $\beta = \alpha \xi$ .

$$\xi = \alpha^{-1}\beta = S\alpha^{-1}\beta + V\alpha^{-1}\beta. \qquad \qquad \therefore \quad S\alpha^{-1}\beta = 0, \quad \xi = V\alpha^{-1}\beta,$$

and  $\xi$  is a constant vector perpendicular to  $\alpha$  and to  $\beta$ , and therefore locates a point.

109. Locus of  $\xi$  in  $V\alpha\xi = \beta$ .

$$\beta \perp \alpha$$
.  $\alpha \xi = S\alpha \xi + V\alpha \xi = x + \beta$ .

$$\therefore \quad \xi = \alpha^{-1}x + \alpha^{-1}\beta = z\alpha + \gamma. \quad [\text{since } \alpha \perp \beta,$$

and the locus of  $\xi$  is a vector through the point determined by  $\gamma(\perp \alpha, \perp \beta)$ , and parallel to  $\alpha$ .

Notice the difference in the reading and interpretation of the equations of this and the previous section. In one  $\beta$  is the whole of  $\alpha \xi$ , in the other only the vector part.

**110.** Locus of  $\xi$  in  $S\alpha\xi=0$ ,  $S\alpha\beta=0$ ,  $S\beta\xi=-c$ .

 $S\alpha\xi=0$  limits  $\xi$  to the plane of  $\alpha$ .

 $S\alpha\beta = 0$  puts  $\beta$  in the same plane.

$$S\beta\xi = -c$$
 or  $bx \cos\theta = c$ ,

where b and x are the tensors of  $\beta$ ,  $\xi$  respectively, makes the projection of  $\xi$  on  $\beta$  a constant, viz.,  $x \cos \theta = \frac{c}{b}$ . Therefore the locus of  $\xi$  is a line  $\perp$  to  $\beta$  in the plane of  $\alpha$  and through the point of  $\beta$  distant from the origin  $\frac{c}{b}$ .

111. From the table of § 85, we find

$$\alpha\beta = \cos \phi + \varepsilon \sin \phi. \qquad (\alpha + \beta)(\alpha + \beta) = (\alpha + \beta)^{2}$$

$$\beta\alpha = \cos \phi - \varepsilon \sin \phi. \qquad = \alpha^{2} + \alpha\beta + \beta\alpha + \beta^{2}.$$

$$\therefore \alpha\beta + \beta\alpha = 2S\alpha\beta = 2S\beta\alpha. \qquad = \alpha^{2} + 2S\alpha\beta + \beta^{2}.$$

$$\alpha\beta - \beta\alpha = 2V\alpha\beta = -2V\beta\alpha.$$

112. By §§ 53, 54, the last equation of § 111 becomes the well-known formula for triangle,

$$c^2 = a^2 + b^2 - 2ab \cos \theta$$
.

113. As in § 111, 
$$(\alpha - \beta)^2 = \alpha^2 - 2S\alpha\beta + \beta^2$$
.  

$$(\alpha +) (\alpha - \beta) = \alpha^2 - \alpha\beta + \beta\alpha - \beta^2$$

$$= \alpha^2 - (\alpha\beta - \beta\alpha) - \beta^2$$

$$= \alpha^2 - 2V\alpha\beta - \beta^2$$
.  

$$\alpha\beta \cdot \beta\alpha = (S\alpha\beta + V\alpha\beta)(S\alpha\beta - V\alpha\beta)$$

$$= (S\alpha\beta)^2 - (V\alpha\beta)^2$$

$$= a^2b^2\cos^2 D - \varepsilon^2\sin^2 D \cdot a^2b^2$$

$$= a^2b^2(\cos^2 D + \sin^2 D)$$

$$= (T\alpha\beta)^2$$
.  

$$\alpha\beta \cdot \beta\alpha = \alpha\beta^2\alpha \qquad \text{by the associative law.}$$

$$= \alpha^2\beta^2 \qquad \text{since } \beta^2 \text{ is a scalar.}$$


$$\alpha\beta \cdot \beta\alpha = \alpha\beta^{2}\alpha \qquad \text{by the associative law}$$

$$= \alpha^{2}\beta^{2} \qquad \text{since } \beta^{2} \text{ is a scalar.}$$

$$\neq (\alpha\beta)^{2}. \qquad \text{Conf. § 88, (5).}$$

$$\alpha\beta \cdot \alpha\beta = (\alpha\beta)^{2} \qquad \alpha\beta \cdot \alpha\beta = \cos^{2}D + \varepsilon 2\sin D \cos D - \sin^{2}D$$
$$= (S+V)^{2}. \qquad = \cos 2D + \varepsilon \sin 2D. \quad \text{Conf. § 98.}$$

114. Applications. If the diagonals of a parallelegram are perpendicular to each other, the parallelogram is a rhombus.



$$S(\alpha + \beta)(\beta - \alpha) = 0 \qquad \S 56.$$
  
=  $\beta^2 - \alpha^2$ .  $\S 113$ .

$$\therefore (T\beta)^2 = (T\alpha)^2. \quad \text{Q.E.D.}$$

115. The joins \* of the mid-points of the sides of a rhombus are at right angles.

$$\gamma = \frac{1}{2}(\alpha + \beta).$$

$$\delta = \frac{1}{2}(\beta - \alpha).$$

$$\gamma \delta = \frac{1}{4}(\alpha\beta - \alpha^2 + \beta^2 - \beta\alpha)$$

$$= \frac{1}{4}(2V\alpha\beta + \beta^2 - \alpha^2).$$
§ 113.

$$\therefore S\gamma \delta = \frac{1}{4}(\beta^2 - \alpha^2) = 0.$$
 [since  $T\alpha = T\beta$ .
$$\therefore \gamma \perp \delta.$$
Q.E.D.


\* Join = the line joining.

116. In any plane triangle to find a side in terms of the other two sides and the opposite angles.

$$\alpha + \beta = \gamma$$
.

Multiplying by (or into)  $\alpha$ , (or any vector, in order to get a quaternion)

$$\alpha^2 + \alpha\beta = \alpha\gamma$$
.



Taking the scalar parts, we have

$$-a^2 + S\alpha\beta = S\alpha\gamma. \qquad -a^2 - ab\cos(180^\circ - C) = -ac\cos B.$$

$$\therefore$$
  $a=b\cos C+c\cos B$ . (Conf. Trigonom.) Q.E.I.

117. Had we taken the vector parts in § 116, we would have found

$$V\alpha\beta = V\alpha\gamma$$
.  $ab \sin C \cdot \varepsilon = ac \sin B \cdot \varepsilon$ .

or  $b \sin C = c \sin B$ . (Conf. Trigonom.)

118. In § 116, had we divided by some vector in order to get a quaternion, say  $\gamma$ , then we would have had the same result,  $c=a\cos B+b\cos A$ .

119. Had we divided by  $\alpha$  or  $\beta$ , say  $\alpha$ , then if the triangle were right-angled at C, since  $S^{\beta}_{\alpha} = 0$ , we would get

$$1 = S\frac{\gamma}{\alpha} = \frac{c}{a} \cos B$$
, or  $\cos B = \frac{a}{c}$ .

120. Had we taken the vector parts in § 118, we should have had  $a \sin B = b \sin A$ . (Law of Sines, Trig.)

Had we taken vector parts in § 119, then since  $\sin C = 1$ ,

$$\sin B = \frac{b}{c}.$$

121.  $\sqrt{i}$  is representable by a prolate spheroidal shell with its major axis i and its two minor axes  $\frac{1}{2}i$  and  $\frac{1}{2}k$ .

Repetitions of this process tend toward the unit shell and  $i^{\circ} = 1$ 

In the same way  $\sqrt{\alpha i} = \sqrt{\alpha} \sqrt{i}$  is representable by a prolate shell whose major axis is i and whose other axes are 1i, 1k, leading similarly to  $(\alpha\beta)^{\circ}=1$ .

From § 113, since,

$$\sqrt{(\alpha\beta)^2} = \sqrt{\cos 2D + \varepsilon \sin 2D} = \cos D + \varepsilon \sin D.$$

$$\therefore (\alpha\beta)^\circ = \cos 0^\circ = 1,$$

the same as above, and agreeing with § 89.

122. Formulæ for reference and practice.

$$\theta$$
 = angle from  $\alpha$  to  $\beta$ .  
 $D$  = angle of the quaternion. [§ 76.  $p, q, r \dots$  = quaternions.

1. 
$$q = Tq(-\cos\theta + \varepsilon\sin\theta) = \alpha\beta$$
.

2. 
$$=Tq(\cos D+\varepsilon\sin D)$$
.

3. = 
$$Tq \cdot Uq = Tq \varepsilon^{\frac{2}{\pi}D}$$
. 4.  $q = Sq + Vq$ .

**5.** 
$$Kq = Sq - Vq = Tq(\cos D - \varepsilon \sin D) = Tq\varepsilon^{-\frac{2}{\pi}D}$$
.

6. 
$$Sq = Tq \cos D = Tq(-\cos \theta)$$
.

7. 
$$TVq = Tq \sin \theta = Tq \sin D$$
.

8. 
$$Vq = TVq \cdot UVq = Tq \sin D \cdot \epsilon$$
.

**9.** 
$$\varepsilon^{\theta+\theta'}$$
. **10.**  $Sq = \frac{1}{2}(q + Kq)$ .

11. 
$$Vq = \frac{1}{2}(q - Kq)$$
.

**12.** 
$$(Tq)^2 = qKq = Kq \cdot q = (Sq)^2 - (Vq)^2$$
.

**13.** 
$$(Tq)^2 = (Sq)^2 + (TVq)^2$$
. **20.**  $T\alpha^2 = -a^2$ .

**14.** 
$$q^{-1} = \frac{Kq}{(Tq)^2}$$
.

**15.** 
$$KKq = q$$
.

16. 
$$Kx = x$$
.

17. 
$$Rq = q^{-1}$$
.

18. 
$$K(-q) = -Kq$$
.

19. 
$$K\alpha = -a\alpha$$
.

**21.** 
$$S\alpha = 0$$
.

**22.** 
$$V\alpha = \alpha$$
.

**23.** 
$$K\alpha\beta = \beta\alpha$$
.

**24.** 
$$S\alpha\beta = S\beta\alpha$$
.

**25.** 
$$V\alpha\beta = -V\beta\alpha$$
.

**26.** 
$$\alpha\beta + \beta\alpha = 2S\alpha\beta$$
.

**27.** 
$$\alpha\beta - \beta\alpha = 2V\alpha\beta$$
.

```
28. (\alpha + \beta)^2 = \alpha^2 + 2S\alpha\beta + \beta^2.
29. T(pqr \dots) = Tp \cdot Tq \cdot Tr \dots
30. U(pqr \dots) = Up \cdot Uq \cdot Ur \dots
31. S(pqr...) = S(qr...p) = S(r...pq) = ...
32. K(pqr \dots) = \dots Kr \cdot Kq \cdot Kp.
33. S(xp+yq+...)*=xSp+ySq+...
34. V(xp + yq + ...) * = xVp + yVq + ...
35. S(q+r+s+...) = Sq+Sr+Ss+...
36. V(q+r+s+...) = Vq + Vr + Vs +...
37. Vq'q = Sq'Vq + SqVq' + V(Vq'Vq).
38. S(\alpha \cdot \beta \gamma) = S\alpha(S\beta \gamma + V\beta \gamma) = S\alpha V\beta \gamma.
                                                                            [M=m+u]
39. S\alpha VM\beta = S\alpha M\beta.
                      =S\alpha(m+\mu)\beta=mS\alpha\beta+S\cdot\alpha\mu\beta.
40.
                      =mS\beta\alpha-S\beta\mu\alpha.
41.
                                                                                                    [§ 51.
                      =S\beta m\alpha - S\beta \mu\alpha = S \cdot \beta KM\alpha.
42.
                                                                                           [§ 122 (5).
43. S(a+\alpha)(b+\beta) = S(b+\beta)(a+\alpha).
44. Spq = Sqp.
45. K\alpha\beta\gamma = K(\alpha\beta\cdot\gamma) = K\gamma\cdot K\alpha\beta = -\gamma\cdot\beta\alpha.
46. \alpha\beta \cdot \gamma - \gamma \cdot \beta\alpha = 2S\alpha\beta\gamma = -2S\gamma\beta\alpha.
47. S\alpha\beta\gamma = S(\alpha\beta\cdot\gamma) = S(p\cdot q) = S(q\cdot p).
48.
                 =S(\gamma\alpha\beta)=S\gamma\alpha\beta=S\beta\gamma\alpha.
49. S(\alpha\beta \cdot \gamma) = S(S\alpha\beta + V\alpha\beta)\gamma.
                      = S \cdot \gamma V \alpha \beta = -S \cdot \gamma V \beta \alpha.
50.
                      = -S \cdot \gamma (V\beta \alpha + S\beta \alpha).
51.
52.
                      =-S\gamma\beta\alpha.
53.
                      =-S\beta\alpha\gamma.
54. S(\alpha_1\alpha_2...\alpha)_n = (-1)^n S(\alpha_n...\alpha_2\alpha_1).
55. S\alpha\beta\gamma = S\alpha V\beta\gamma.
                 =S \cdot \alpha V \beta \gamma = S \cdot \beta V \gamma \alpha = S \gamma V \alpha \beta.
56.
57. \alpha\beta\gamma + \gamma\beta\alpha = 2V\alpha\beta\gamma = 2V\gamma\beta\alpha.
58. 2V\alpha\beta\gamma = \alpha\beta\gamma + (\alpha\gamma\beta - \alpha\gamma\beta - \gamma\alpha\beta + \gamma\alpha\beta) + \gamma\beta\alpha.
                                                                                                       [57.
```

<sup>\*</sup> See Appendix.

<ul> <li>1 Τι έχ = ε έν - νε - εν - γε έ - γι ε - βαλ.</li> <li>2 ε ν έγ - άναν - γναε .</li> <li>3 Τι έν - ανέν - Γιανέν.</li> <li>4 = Γα έν - βεν.</li> <li>4 = Για Γέχ.</li> </ul>	<b>]26</b> .
43. $ = V \cdot a V \hat{r} r $ 44. $ = -\tilde{a} S \cdot a + r S a \tilde{r} $	<b>[60.</b>
$=-\Gamma\cdot\Gamma\stackrel{\circ}{\sim}\alpha.$	[62, 25.
$\mathfrak{M}_{n,n} = T \cdot T \cdot \tilde{p} = \tilde{p} \cdot \tilde{p} = $	[ <del>64</del> .
#7. T Tu? V-1 = -78. Vai +18. Va? 7.	[ <b>63</b> , <b>64</b> .
$=-\gamma SaSi-iSaS$	[56, 24.
$60  T  Ta?  Tr? = -aS  \hat{z} \text{ Tr} \hat{z} - \hat{z} S  \text{ Tr} \hat{z} \text{ on}$	[ <b>66</b> .
$\mathbf{T} \mathbf{I}. \qquad = \mathbf{a} \hat{\mathbf{S}} \hat{\mathbf{c}} \cdot \hat{\mathbf{c}} - \hat{\mathbf{p}} \hat{\mathbf{S}} \mathbf{a} \cdot \hat{\mathbf{c}}.$	<b>[56</b> .
72 i Suir=uSiri+jSrai+ySail.	[68, 70.
72. T Taility = 78 Tai i - 58 7 Tai.	[67.
73. $= r \hat{S} \hat{z} \hat{T} \hat{a} \hat{z} + \hat{z} \hat{S} \hat{z} \hat{z}.$	[24, 56.
$74. = 55527. \qquad ISFVa$	$\dot{z} = 0$ , 60.
75. abasy = V · Vsa Vay.	[74.
76. $\Delta S \Delta S \gamma = \Gamma \cdot V \gamma S \Gamma \Delta \Delta$ .	[48, 75.
77. $\tau S_2 \hat{z}_{\tau} = \Gamma \cdot \Gamma_2 \cdot \Gamma_{\tau} \hat{z}_{\tau}$	
78. i =S.i - [S]i - kSci .	[71.

122. In 63 , 64 ,  $V \cdot \alpha V_{eff}$  is perpendicular to  $\alpha$  and coplanar with  $\beta$ ,  $\gamma$ . That it is perpendicular to  $\alpha$  can be shown by multiplying by  $\alpha$  and taking scalars thus,

 $S_2 \left[ V \cdot a V_{F_2} \right] = S_2 \cdot S_2 \cdot F - S_2 \cdot S_2 \cdot F = 0.$  [63, 64.

It is in the plane of Fr since

$$V \cdot \alpha V_{FT}^2 = 7S\alpha \hat{F} - \hat{\mu} S\alpha \hat{T}$$
. [63, 64.

124. In 58, 60 Vaff is of the form xa + yb + xy and is therefore the intermediate disgonal of the parallelophped of which the edges are a  $Sfy_1 + bSay_2 + ySab_2$ .

**125.** In (67), since 
$$S \cdot \gamma S \alpha \beta = S \cdot \delta S \alpha \beta = 0$$
,

$$\begin{split} V \cdot V \alpha \beta V \gamma \delta &= \delta S \{ S \alpha \beta \cdot \gamma + V \alpha \beta \cdot \gamma \} - \gamma S \{ \delta S \alpha \beta + \delta V \alpha \beta \} \\ &= \delta S (S \alpha \beta + V \alpha \beta) \gamma - \gamma S \delta (S \alpha \beta + V \alpha \beta) \\ &= \delta S \alpha \beta \gamma - \gamma S \delta \alpha \beta, \end{split}$$

and it is evident that  $V \cdot V \alpha \beta V \gamma \delta$  is coplanar with  $\delta$  and  $\gamma$ . Moreover, since

$$V \cdot V \alpha \beta V \gamma \delta = V \cdot V \delta \gamma V \alpha \beta = \beta S \delta \gamma \alpha - \alpha S \beta \delta \gamma$$
, [25, 52, 67.

it is also coplanar with  $\alpha$  and  $\beta$ , and therefore must be along the intersection of the planes of  $\alpha$ ,  $\beta$  and  $\gamma$ ,  $\delta$ .

126. In (71),  $\delta S\alpha\beta\gamma$  is the intermediate diagonal of the parallelopiped of which the three edges are  $\alpha S\beta\gamma\delta$ ,  $\beta S\gamma\alpha\delta$ ,  $\gamma S\alpha\beta\delta$ .

#### CHAPTER XI

### QUATERNION EQUATIONS OF THE FIRST DEGREE

127. An equation of the first degree with respect to an unknown quaternion X, is that which contains the quaternion to the first power together with known quaternions, either isolated or under the symbols S or V.

The general equation will then have the form

$$\Sigma AXB + \Sigma C \cdot SAXB + \Sigma D \cdot VAXB \cdot E = F.$$

The third term assumes the form of the first two if we replace VAXB by AXB-SAXB, so that the general equation reduces to the form

$$\Sigma AXB + \Sigma C \cdot SAXB = F$$
.

128. To resolve this equation we decompose the quaternions into their scalar and vector parts. Thus, putting  $A = a + \alpha$ ,  $B = b + \beta$ , etc., we have

$$\sum (a+\alpha)(x+\xi)(b+\beta) + \sum (c+\gamma)S \cdot (a+\alpha)(x+\xi)(b+\beta) = d+\delta.$$

The sum of the scalar parts of this will be found to be embraced in the general term  $S\alpha\xi$ , therefore the scalar part of the equation is  $S\alpha\xi = d$ .


129. Solution of the linear scalar equation  $Sa\xi = d$ . This may be written, § 122, (33),

$$S\alpha(\xi-d\alpha^{-1})=0,$$
  $[\alpha d\alpha^{-1}=d.$ 

where, § 56, (2), evidently  $(\xi - d\alpha^{-1})$  is some vector  $\beta$ ,  $\perp \alpha$ , or  $\beta = V\alpha \gamma$ , where  $\gamma$  is an arbitrary vector. Therefore,

$$\xi = d\alpha^{-1} + \beta = d\alpha^{-1} + V\alpha r.$$

The geometrical interpretation of this is shown in the diagram, where since  $\gamma$  is arbitrary, the locus of the extremity of  $\xi$  must be the plane  $\perp \alpha$  and through the point  $d\alpha^{-1}$ .



$$abx\gamma + aS\xi\beta \cdot \gamma + xS\alpha\beta \cdot \gamma + bS\alpha\xi + S\alpha\xi\beta \cdot \gamma$$
.

Multiplying out the first term we find the vector part to be

$$ax\beta + ab\xi + a\xi\beta + bx\alpha + x\alpha\beta + b\alpha\xi + \alpha\xi\beta$$
.

Of these forms

$$a\xi\beta = a(S\xi\beta + V\xi\beta),$$
  $b\alpha\xi = b(S\alpha\xi + V\alpha\xi),$   $x\alpha\beta = x(S\alpha\beta + V\alpha\beta),$   $\alpha\xi\beta = S\alpha\xi\beta + V\alpha\xi\beta$ 

$$= S\alpha\xi\beta + \alpha S\xi\beta - \xi S\alpha\beta + \beta S\alpha\xi. \quad [\S 122, (61), (64)].$$

Of the final forms,  $abx \cdot \gamma$ ,  $aS\xi\beta \cdot \gamma$ ,  $xS\alpha\beta \cdot \gamma$ ,  $b \cdot S\alpha\xi \cdot \gamma$ ,  $S\alpha\xi\beta \cdot \gamma$ ,  $ax\beta$ ,  $bx\alpha$ ,  $aS\xi\beta$ ,  $xV\alpha\beta$ ,  $bS\alpha\xi$ , are comprehended under the general form,

$$\alpha S\beta \xi$$
;

 $ab\xi$ ,  $\xi S\alpha\beta$  are comprehended under the general form,

$$Vm\xi = V(ab + S\alpha\beta + ...);$$

<sup>\*</sup> Or neglected as purely scalar.

 $aV\xi\beta$ ,  $bV\alpha\xi$  are comprehended under the form,

$$V \mu \xi = (V \beta \xi + V \alpha \xi + V \gamma \xi + \dots).$$

Hence the vector part of the general equation becomes

$$\Sigma \alpha S \beta \xi + V m \xi + V \mu \xi = \delta,$$
 or  $\Sigma \alpha S \beta \xi + V (m + \mu) \xi = \delta.$  [§ 122, (34). or  $\Sigma \alpha S \beta \xi + V \cdot M \xi = \delta.$  [§ 61.

This is generally abbreviated under the functional sign,

$$\sum \alpha S\beta \xi + V \cdot M\xi = \phi \xi = \delta.$$

131. Hamilton called this a linear vector function of the vector  $\xi$ . Considered as an operator its application to  $\xi$  has many interesting results, some of which will be investigated in the following pages.

We will consider first the properties of  $\phi$  itself, and then the result of its application to the vector  $\xi$ .

For reasons given later this function is called a strain function.

132. Properties of  $\phi$ . Since § 122 (35),

$$\beta S\alpha(\xi + \eta + \dots) = \beta S\alpha\xi + \beta S\alpha\eta + \dots,$$

and § 122, (36),

$$V \cdot M(\xi + \eta + \dots) = V \cdot M\xi + V \cdot M\eta + \dots;$$

$$\therefore \quad \Sigma \beta S \alpha(\xi + \eta + \dots) + V \cdot M(\xi + \eta + \dots)$$

$$= (\Sigma \beta S \alpha \xi + V \cdot M\xi) + (\Sigma \beta S \alpha \eta + V \cdot M\eta) + \dots,$$
or
$$\phi(\xi + \eta + \dots) = \phi \xi + \phi \eta + \dots,$$

that is,  $\phi$  is distributive over a sum.

133. If  $\xi = \eta = \dots$  to n terms,

then

$$\phi n\xi = n\phi \xi$$
,

that is,  $\phi$  is commutative with a scalar factor.

**134.** If we define  $\phi^{-1}$  by the equation,  $\phi^{-1}\phi = 1, *$  then

$$\phi^{-1}\phi\xi = \phi^{-1}\delta = \xi,$$

and from § 132,

$$\phi^{-1}\phi(\xi+\eta+\ldots)=\phi^{-1}(\phi\xi+\phi\eta+\ldots),$$
  
$$\xi+\eta+\ldots=\phi^{-1}(\phi\xi+\phi\eta+\ldots).$$

or

But  $\phi \xi = \delta$ , etc., and  $\phi^{-1} \delta = \xi$ , etc., and therefore,

$$\phi^{-1}\delta + \phi^{-1}\delta_1 + \ldots = \phi^{-1}(\delta + \delta_1 + \ldots).$$

Hence  $\phi^{-1}$  has the same properties as  $\phi$ .

By repeated applications we can easily get, for all integral values of k,

$$\phi^k \phi^{-k} \xi = \xi$$
.

135. Conjugate strain functions. Operating on  $\phi \xi = \sum \alpha S \beta \xi + V \cdot M \xi$  with  $S \cdot \eta$ , we get

$$\begin{split} S \cdot \eta \phi \xi &= S \eta ( \sum \alpha S \beta \xi + V \cdot M \xi ) \\ &= \sum S \cdot \eta \alpha S \beta \xi + S \cdot \eta V \cdot M \xi \\ &= \sum S \xi \beta S \alpha \eta + S \eta (m + \mu) \xi \\ &= \sum S \xi \beta S \alpha \eta + S \xi (m + \mu) \eta \quad \text{[§ 122, (24), (38).} \\ &= \sum S \xi \beta S \alpha \eta + S \xi (m - \mu) \eta \quad \text{[§ 122 (52).} \\ &= S \xi ( \sum \beta S \alpha \eta + V K M \eta ) \quad \text{[§ 122 (39), (5).} \\ &= S \xi \phi' \eta. \end{split}$$

 $\phi$  and  $\phi'$  are called **conjugate strain functions.** They evidently differ in the interchange of the known vector  $\alpha$ ,  $\beta$ , and of the quaternion M and its conjugate KM.

<sup>\*</sup> See Appendix.

136. When  $\phi = \phi'$ , that is, the function conjugates into itself, the functions are called self-conjugate strain functions, in which case  $S\eta\phi\xi = S\xi\phi\eta$ .

137. Types of self-conjugate functions. Since

$$S\xi\phi\eta = S\eta\phi'\xi$$
.  $\therefore S\eta\phi\xi = S\xi\phi'\eta$ ,

and writing \*  $(\phi + \phi')\xi$  for  $\phi\xi + \phi'\xi$ ,

$$S\eta(\phi+\phi')\xi=S\xi(\phi+\phi')\eta$$

which shows that the operator  $\phi + \phi'$  is always self conjugate.

138. Furthermore,  $S\xi\phi\xi = S\xi\phi'\xi$ ,

whence

$$S\xi(\phi-\phi')\xi=0$$
,

and therefore

the vector  $(\phi - \phi')\xi$  is perpendicular to  $\xi$ 

or

$$(\phi - \phi')\xi = V\varepsilon\xi$$
,

where  $\epsilon$  is some unknown vector.

Consequently, 
$$\phi \xi = \frac{1}{2}(\phi + \phi')\xi + \frac{1}{2}(\phi - \phi')\xi$$
  
=  $\frac{1}{2}(\phi + \phi)'\xi + \frac{1}{2}V\varepsilon\xi$ ,

which shows that a linear vector function of  $\xi$  differs from a self-conjugate function only by a term of the form  $V \in \xi$ . If it is already self conjugate the vector  $\varepsilon = 0$ .

**139.** Since

$$\begin{split} \mathcal{S} \cdot \xi \phi \phi' \eta = & S \xi \phi(\phi' \eta) & S \cdot \xi \phi \phi' \eta = & S \cdot \eta \phi(\phi' \xi) & \text{[§ 135.} \\ &= & S \cdot \phi' \eta \phi' \xi & \text{[§ 135.} & = & S \eta \phi \phi' \xi, \\ &= & S \cdot \phi' \xi \phi' \eta; & \text{[§ 122 (24).} \end{split}$$

therefore  $\phi\phi'$  is a self-conjugate strain function.

<sup>\*</sup> This is allowable because the functional symbol  $\phi$  has the same properties as an algebraic factor (distributive over a sum and commutative with scalar factors) and can be treated like one. See Appendix.

**140.** If by  $(\phi + g)\xi$  we understand  $\phi \xi + g\xi$ , where g is a scalar,  $\phi + g$  is also a linear vector function, since it has all the properties of  $\phi$  as the reader can easily demonstrate for himself. Hence

$$S \cdot \xi(\phi + g) \eta = S \cdot \xi(\phi \eta + g \eta) = S \cdot \xi \phi \eta + S \cdot \xi g \eta,$$

$$= S \cdot \eta \phi' \xi + g S \cdot \eta \xi = S \cdot \eta(\phi' + g) \xi,$$

or  $\phi' + g$  is a conjugate function to  $\phi + g$ .

141. Application of  $\phi$  to a vector  $\xi$ . If an elastic solid. that is, the vector connecting the several elements, be

subjected to the operation  $\phi$ , then all its particles, for instance, those determined by the vectors  $\alpha$ ,  $\beta$ ,  $\gamma$ , are displaced to positions determined by the vectors  $\phi \alpha$ ,  $\phi \beta$ ,  $\phi \gamma$ . In general, any particle whose vector is  $\xi$  occupies



after the operation the position whose vector is  $\phi \xi$ .

Also any vector  $\alpha$  is displaced to the position  $\phi \alpha$ , for since

$$\alpha = \gamma - \beta,$$

$$\phi \alpha = \phi \gamma - \phi \beta.$$

Hence, Any straight line of particles parallel to  $\alpha$ , say  $x\alpha$ , is homogeneously stretched and turned by the operation  $\phi$  into a straight line of particles parallel to  $\phi \alpha$ , and the ratio of extension is  $\frac{x\phi\alpha}{r\alpha} = \frac{\phi\alpha}{\alpha}$ .

The operation  $\phi$  is called a strain and the property that parallel lengths are strained into parallel lengths and stretched proportionally is the physical definition of linear homogeneous strain. Portions of the body originally equal, similar and similarly placed remain after the strain equal, similar and similarly placed.

142. If in the general equation of the first degree with respect to an unknown quaternion, § 127, instead of separating the scalar and vector parts, we had merely indicated the vector part, thus

$$\Sigma VQXR = VF$$
.

this operator VQ()R must of course be the undeveloped form of  $\phi$ .

If  $R = Q^{-1}$ , we have the *rotator* of § 102 as a special case of  $\phi$ .

If Q and R degrade into scalars, we have  $\Sigma VmX = nX$ , or dilatation merely.

A combination of rotation and dilatation makes the strain just defined.

143. Properties of  $\phi$ . To get a slightly different view, let us suppose  $\alpha$ ,  $\beta$ ,  $\gamma$  to be unit vectors at right angles to each other. Hence, § 122 (78),

$$\xi = -(\alpha S\alpha \xi + \beta S\beta \xi + \gamma S\gamma \xi).$$

By the definition of homogeneous strain this is changed into

$$\xi' = \phi \xi = -(\alpha' S \alpha \xi + \beta' S \beta \xi + \gamma' S \gamma \xi),$$

where  $\alpha'$ ,  $\beta'$ ,  $\gamma'$  are three vectors upon which the same proportional distances  $S\alpha\xi$ ,  $S\beta\xi$ ,  $S\gamma\xi$  are laid off. This is necessary in order to preserve the similarity required by homogeneous strain. By substituting  $\alpha$ ,  $\beta$ ,  $\gamma$  for  $\xi$ , we find

$$\alpha' = \phi \alpha$$
,  $\beta' = \phi \beta$ ,  $\gamma' = \phi \gamma$ .

$$\begin{split} &\phi\alpha=\alpha'=-\left(\alpha S\alpha\alpha'+\beta S\beta\alpha'+\gamma S\gamma\alpha'\right)=\alpha A_{\alpha}+\beta B_{\alpha}+\gamma C_{\alpha},\\ &\phi\beta=\beta'=-\left(\alpha S\alpha\beta'+\beta S\beta\beta'+\gamma S\gamma\beta'\right)=\alpha A_{\beta}+\beta B_{\beta}+\gamma C_{\beta},\\ &\phi\gamma=\gamma'=-\left(\alpha S\alpha\gamma'+\beta S\beta\gamma'+\gamma S\gamma\gamma'\right)=\alpha A_{\tau}+\beta B_{\tau}+\gamma C_{\tau}, \end{split}$$

which equations contain *nine* arbitrary constants,  $S\alpha\alpha'$ ,  $S\beta\alpha'$ , etc., since  $\alpha'$ ,  $\beta'$ ,  $\gamma'$  are entirely independent of  $\alpha$ ,  $\beta$ ,  $\gamma$ .

Operating on  $\phi \xi$  with  $S \cdot \eta$  we have

$$S \cdot \eta \phi \xi = -\left(S\alpha' \eta S\alpha \xi + S\beta' \eta S\beta \xi + S\gamma' \eta S\gamma \xi\right)$$

$$= -S \cdot \xi (\alpha S\alpha' \eta + \beta S\beta' \eta + \gamma S\gamma' \eta), \qquad [\S 122 (24).$$

where the expression in the parenthesis is a linear vector function of  $\eta$ , which will be shown (§ 144) to depend upon the same nine scalars  $A_{\alpha}$ ,  $B_{\alpha}$ , etc., as those in  $\phi$ , and which we may therefore appropriately designate by  $\phi'$ , thus

$$\phi'\eta = -(\alpha S\alpha'\eta + \beta S\beta'\eta + \gamma S\gamma'\eta),$$

whence obviously  $S \cdot \eta \phi \xi = S \cdot \xi \phi' \eta$ , as before.

144. Substituting  $\alpha$ ,  $\beta$ ,  $\gamma$  for  $\eta$  in the expression for  $\phi'\eta$ , we find

$$\phi'\alpha = -(\alpha S\alpha'\alpha + \beta S\beta'\alpha + \gamma S\gamma'\alpha);$$
  

$$\phi'\beta = -(\alpha S\alpha'\beta + \beta S\beta'\beta + \gamma S\gamma'\beta);$$
  

$$\phi'\gamma = -(\alpha S\alpha'\gamma + \beta S\beta'\gamma + \gamma S\gamma'\gamma),$$

or using the notation of § 143,

$$\phi'\alpha = \alpha A_{\alpha} + \beta A_{\beta} + \gamma A_{\gamma};$$
  
$$\phi'\beta = \alpha B_{\alpha} + \beta B_{\beta} + \gamma B_{\gamma};$$
  
$$\phi'\gamma = \alpha C_{\alpha} + \beta C_{\beta} + \gamma C_{\gamma};$$

which shows that  $\phi'$  depends upon the same nine scalars as  $\phi$ .

**145.** If  $\alpha$ ,  $\beta$ ,  $\gamma$  are given non-coplanar vectors, then § 29,

$$\xi = x\alpha + y\beta + z\gamma,$$

whence

$$\phi \xi = x \phi \alpha + y \phi \beta + z \phi \gamma.$$

The vector  $\phi\xi$  is known when the three vectors  $\phi\alpha$ ,  $\phi\beta$ ,  $\phi\gamma$ , are known. Each of these vectors involves three scalar constants as in the case of  $\xi$ , multiples of the reference vectors  $\alpha$ ,  $\beta$ ,  $\gamma$ . Therefore the value of  $\phi$  depends upon nine scalar constants. It has therefore been called a **nonion.** 

#### CHAPTER XII

### APPLICATIONS OF $\phi$

146. Changes of volume due to  $\phi$ .  $\rho$ ,  $\phi \rho$ ,  $\phi^2 \rho$  are, in general, not in one plane, and hence, § 29,

$$\phi^{3}\rho = m_{2}\phi^{2}\rho - m_{1}\phi\rho + m\rho$$
, . . . (1)

where  $m_2$ ,  $m_1$ , m are scalars independent of  $\rho$ . The independence is obvious, since we may put  $\alpha$ ,  $\beta$ ,  $\gamma$  in succession for  $\rho$  and thus obtain three equations from which they can be obtained.

From § 59 the volume of the parallelopiped whose three conterminous edges are  $\rho$ ,  $\phi \rho$ ,  $\phi^2 \rho$  is

$$-S \cdot \rho \phi \rho \phi^2 \rho$$
.

After the strain this volume is

$$-S \cdot \phi \rho \phi^2 \rho \phi^3 \rho$$
.

Multiplying Eq. (1) by  $\phi\rho\phi^2\rho$  and dividing by  $\rho\phi\rho\phi^2\rho$ , we have

$$\frac{S \cdot \phi \rho \phi^2 \rho \phi^3 \rho}{S \cdot \rho \phi \rho \phi^2 \rho} = m. \qquad (2)$$

This ratio of dilatation m due to  $\phi$ , the ratio between the volumes after and before the strain, is called the modulus of the strain  $\phi$ .

147. By referring to the equations of § 143 we see that the scalar part of the product  $\phi \alpha \phi \beta \phi \gamma$  is confined to those terms in which all three of the vectors  $\alpha$ ,  $\beta$ ,  $\gamma$  appear, that is, taking  $S \cdot \alpha \beta \gamma = 1$ ,

$$\begin{split} m = & \frac{S \cdot \phi \alpha \phi \beta \phi \gamma}{S \cdot \alpha \beta \gamma} \\ = & A_{\alpha} B_{\beta} C_{\gamma} - A_{\alpha} B_{\gamma} C_{\beta} - B_{\alpha} A_{\beta} C_{\gamma} + B_{\alpha} A_{\gamma} C_{\beta} + \\ & + C_{\alpha} (A_{\beta} B_{\gamma} - A_{\gamma} B_{\beta}). \end{split}$$

In the same way from § 144,

$$\frac{S \cdot \phi' \alpha \phi' \beta \phi' \gamma}{S \cdot \alpha \beta \gamma} = A_{\alpha} B_{\beta} C_{\gamma} - A_{\alpha} B_{\gamma} C_{\beta} - \text{, etc.}$$

$$= m.$$

Hence, Conjugate strains produce equal changes of volume. 148. Special values of m. If m=1 there is no change of volume caused by the strain.

If m=0, there results a zero volume due to the solid becoming strained into a plane, a line, or a point, and in this case  $\phi$  is called a **null function**, singly, doubly, or triply **null** according as the strain results in a plane, a line or a point. The plane, line, or point into which the solid is strained is called the **strain plane** of  $\phi$ , strain line, etc.

When m=0, then  $S\phi\alpha\phi\beta\phi\gamma$ , § 147, becomes zero and, § 60,  $\phi\alpha$ ,  $\phi\beta$ ,  $\phi\gamma$  are coplanar vectors, and we can have a relation,

$$x\phi\alpha + y\phi\beta + z\phi\gamma = 0,$$
 or § 132, 
$$\phi(x\alpha + y\beta + z\gamma) = 0.$$

The vector,  $x\alpha + y\beta + z\gamma$ , whose strain, i.e., the result of the application of the strain function  $\phi$ , is zero is called a **null direction** of  $\phi$ . In this case  $\phi$  is **singly null**, unless  $\phi\alpha$ ,  $\phi\beta$ ,  $\phi\gamma$  becomes collinear.

149. If there is only one vector, say  $\alpha$ , whose strain is zero, i.e.,  $\phi \alpha = 0$ , then

$$S \cdot \phi \alpha \phi \beta \phi \gamma = 0$$
,

and therefore m=0, and  $\phi\beta$  cannot be parallel to  $\phi\gamma$  (i.e.,  $\phi\beta \neq x\phi\gamma$ ), for otherwise  $\phi(\beta-x\gamma)$  would be equal to zero and there would be a second vector,  $\beta-x\gamma$ , with a strain of zero, which is contrary to the hypothesis. Hence, if there is only one null vector and  $\rho = x\alpha + y\beta + z\gamma$ , then

$$\phi \rho = y \phi \beta + z \phi \gamma$$

and  $\phi \rho$  is a vector in a plane parallel to  $\phi \beta$ ,  $\phi \gamma$ , that is,  $\phi$  is singly null.

**150.** If there are two (only) vectors whose strain is zero, say  $\phi \alpha = 0$ ,  $\phi \beta = 0$ , then  $\phi \rho = z \phi \gamma$ , which confines  $\phi \rho$ , the strain of the general vector  $\rho$ , to the line parallel to  $\phi \gamma$ , and in this case  $\phi$  is doubly null.

**151.** Similarly, if  $\phi \alpha = 0$ ,  $\phi \beta = 0$ ,  $\phi \gamma = 0$ , then  $\phi \rho = 0$  for all values of  $\rho$  and  $\phi$  is a triply-null function.

152. When  $\phi$  is singly null, say  $\phi \alpha = 0$ , then the general vector of any point in the strain plane of  $\phi$  is, § 149,

$$\phi \rho = y \phi \beta + z \phi \gamma$$

and all particles that strain into this plane have the vectors,

$$\rho = x\alpha + y\beta + z\gamma,$$

where x is arbitrary, since  $\phi \alpha = 0$ , and therefore the locus of  $\rho$  is a line parallel to  $\alpha$ . Hence,

A singly null function strains each of its null lines into a definite point of its strain plane. Similarly,

A doubly-null funcion strains each of its null planes into a definite point of its strain line.

153. Special applications of  $\phi$ . What vectors, if any, are unchanged in position by the strain  $\phi$ ?

If  $\rho$  is unchanged by the strain, then

$$\phi \rho = g\rho$$
,  $\phi^2 \rho = g^2 \rho$ ,  $\phi^3 \rho = g^3 \rho$ ,

and the equation of § 146 becomes

$$(g^3-m_2g^2+m_1g-m)\rho=0$$
, . . . (1)

which must have one real root  $g_1$  and may have three. In other words, the curious fact that however a body may be homogeneously strained, there is always at least one vector whose direction remains unchanged. Hence

$$\phi \rho - g_1 \rho = 0$$
,

and

$$S \cdot \mu \phi \rho - g_1 S \mu \rho = 0$$
,  $\mu = \text{any vector.}$ 

and

$$S \cdot \rho(\phi' \mu - g_1 \mu) = 0 = S \cdot \rho(\phi' - g) \mu.$$
 [§ 135.

Hence, § 56 (2),  $(\phi'-g_1)\mu$  is perpendicular to  $\rho$ .

From this it appears that the operator  $(\phi'-g)$  applied to any vector  $\mu$  throws it into a plane perpendicular to  $\rho$ , or, in other words, cuts off the component of the strained  $\mu$  which is parallel to  $\rho$ .

Also,  $\rho \| V \cdot (\phi' - g_1) \mu(\phi' - g_1) \nu$ ,  $[\mu, \nu]$  any vectors.

or multiplying out

$$\rho \|V(\phi'\mu\phi'\nu - g_1(\phi'\mu\cdot\nu + \mu\phi'\nu) + g_1^2\mu\nu)$$

$$\|V\phi'\mu\phi'\nu - g_1(V\phi'\mu\cdot\nu + V\mu\phi'\nu) + g_1^2V\mu\nu.$$

154. Properties of  $(\phi'-g)$ . Like  $\phi$ , this operator produces a homogeneous strain, as can be shown in a manner similar to that in the case of  $\phi$ . It has the same general properties and has a modulus  $m_g$ . Omitting the primes,

which will not affect the discussion, we have, after expanding.

$$\begin{split} m_{g} = & \frac{S \cdot (\phi - g)\alpha(\phi - g)\beta(\phi - g)\gamma}{S \cdot \alpha\beta\gamma} \\ = & m - g \frac{S(\alpha\phi\beta\phi\gamma + \beta\phi\gamma\phi\alpha + \gamma\phi\alpha\phi\beta)}{S \cdot \alpha\beta\gamma} \\ & + g^{2} \frac{S(\alpha\beta\phi\gamma + \gamma\alpha\phi\beta + \beta\gamma\phi\alpha)}{S \cdot \alpha\beta\gamma} - g^{3} \\ = & m - m'_{1}g + m'_{2}g^{2} - g^{3}. \end{split}$$

155. In § 153 we found the expression.

$$V\cdot(\phi'-g_1)\mu(\phi'-g_1)\nu$$

the vector part of the product of two strained vectors. To investigate this we return to the last equation of § 146, which can be written,

$$m = \frac{S \cdot \phi \alpha \phi \beta \phi \gamma}{S \cdot \alpha \beta \gamma} = \frac{S \cdot (\phi \beta \phi \gamma) \phi \alpha}{S \cdot \alpha \beta \gamma} = \frac{S \alpha \phi' V (\phi \beta \phi \gamma)}{S \cdot \alpha V \beta \gamma}, \quad [\S 135.$$

or since  $\alpha$  is any vector whatever,

$$\phi'V(\phi\beta\phi\gamma) = mV\beta\gamma, \quad . \quad . \quad . \quad . \quad (1)$$

or since, § 147,  $\phi$  and  $\phi'$  have the same modulus,

$$\phi V(\phi'\beta\phi'\gamma)=mV\beta\gamma,$$

Using  $(\phi - g)$  in place of  $\phi$ , we have

$$\begin{split} (\phi - g) V(\phi' - g) \beta(\phi' - g) \gamma &= m_g V \beta \gamma \\ &= (\phi - g) V[\phi' \beta \phi' \gamma - g(\phi' \beta \cdot \gamma + \beta \phi' \gamma) + g^2 \beta \gamma] \\ &= (\phi - g) [V \phi' \beta \phi' \gamma - g(V \phi' \beta \cdot \gamma + V \beta \phi' \gamma) + g^2 V \beta \gamma] \\ &= (\phi - g) [m \phi^{-1} V \beta \gamma - g(V \phi' \beta \cdot \gamma + V \beta \phi' \gamma) + g^2 V \beta \gamma] [(1) \\ &= (m - m'_1 g + m'_2 g^2 - g^3) V \beta \gamma. \end{split}$$

**156.** In this equation and the last expression of § 153, we have the four vectors  $V\beta\gamma$ ,  $V\cdot\phi'\beta\cdot\gamma$ ,  $V\cdot\beta\phi'\gamma$ ,  $\phi V\beta\gamma$ , and we can write

$$\phi V \beta \gamma = x V \beta \gamma + y V \cdot \phi' \beta \cdot \gamma + z V \beta \phi' \gamma.$$

Operate successively with  $S \cdot \alpha$ ,  $S \cdot \beta$ ,  $S \cdot \gamma$  and we get

$$S\beta\gamma\phi'\alpha = xS \cdot \alpha\beta\gamma + yS \cdot \gamma\alpha\phi'\beta + zS \cdot \alpha\beta\phi'\gamma.$$

$$S\beta\gamma\phi'\beta = yS \cdot \gamma\beta\phi'\beta.$$

$$S\beta\gamma\phi'\gamma = zS \cdot \gamma\beta\phi'\gamma.$$

But § 122 (53), y = -1, z = -1, and

$$x = \frac{S \cdot \beta \gamma \phi' \alpha + S \cdot \gamma \alpha \phi' \beta + S \cdot \alpha \beta \phi' \gamma}{S \cdot \alpha \beta \gamma} = m'_{2}.$$
 [§ 154.

157. Hence, § 156,

$$\phi V \beta \gamma = m'_{2} V \beta \gamma - V \phi' \beta \cdot \gamma - V \beta \phi' \gamma,$$

or substituting in the equation of § 155 and using  $\xi$  for  $V\beta\gamma$ , we get

$$(\phi - g)[m\phi^{-1} - g(m'_2 - \phi) + g^2]\xi = (m - m'_1g + m'_2g^2 - g^3)\xi,$$

or expanding,

$$\begin{split} (m+g\phi^2+m'_2g\phi+g^2\phi-gm\phi^{-1}+g^2m'_2-g^2\phi-g^3)\,\xi\\ &=(m-m'_1g+m'_2g^2-g^3)\,\xi, \end{split}$$

or 
$$(\phi^2 - m'_2 \phi - m \phi^{-1}) \xi = -m'_1 \xi$$
, . . . (1)  
or  $(\phi^3 - m'_2 \phi^2 + m'_1 \phi - m) \xi = 0$ ,

where  $\xi$  is any vector whatever.

158. Comparing this with Eq. (1), § 146, we have

$$\begin{split} m_2 = m'_2 &= \frac{S \cdot \alpha\beta\phi\gamma + S \cdot \gamma\alpha\phi\beta + S \cdot \beta\gamma\phi\alpha}{S\alpha\beta\gamma}, \\ m_1 = m'_1 &= \frac{S \cdot \alpha\phi\beta\phi\gamma + S \cdot \beta\phi\gamma\phi\alpha + S \cdot \gamma\phi\alpha\phi\beta}{S\alpha\beta\gamma} \end{split}$$

The expression of § 153 now becomes, by §§ 155, 157, and the equation above,

$$\rho \| (m\phi^{-1} - g_1(m_2 - \phi) + g_1^2) \xi \| \left( \frac{m}{g_1} - (m_2 - g_1)\phi + \phi^2 \right) \xi, ,$$

[dividing by  $g_1$  and operating with  $\phi$  or, multiplying and dividing by  $(\phi - g_1)$  and then substituting the value of

$$\frac{\phi m}{g_1} + m_2 g_1 \phi - g_1^2 \phi$$
,

from (1) of § 153.

$$\rho \| \frac{\phi^3 - m_2 \phi^2 + m\phi - m}{\phi - q_1} \xi. \qquad [\xi = \text{any vector.}]$$

**159.** Since (1) of § 153 is true for all vectors, and since  $\phi$ is commutative with scalars, it can be written,

$$(\phi - g_1)(\phi - g_2)(\phi - g_3)\xi$$
  
 $\rho || (\phi - g_2)(\phi - g_3)\xi.$ 

and, § 158,

that is, the operator  $(\phi - g_2)(\phi - g_3)$  when applied to any vector whatever results in a vector parallel to  $\rho$ , where

$$(\phi - g_1)\rho = 0,$$

or

$$\phi \rho = g_1 \rho$$

 $\rho$  being unchanged in direction by the strain  $\phi$ .

Now

160. We can fortify this by another line of attack. As before, § 153, under the condition,  $\phi \xi = g \xi$ , we have

$$(g^3 - m_2 g^2 + m_1 g - m) \xi = 0.$$

Supposing for the moment that the three roots are real, the solution of the problem will be given by one of the directions  $\alpha$ ,  $\beta$ ,  $\gamma$  which satisfies the conditions,

$$(\phi - g_1)\alpha = 0$$
,  $(\phi - g_2)\beta = 0$ ,  $(\phi - g_3)\beta = 0$ . (1)  
 $\xi = a\alpha + b\beta + c\gamma$ .

Operating with  $(\dot{\phi}-g_1)$ , we get

$$\begin{split} (\phi-g_1)\xi = &b(\phi-g_1)\beta + c(\phi-g_1)\gamma \\ = &(b\phi-bg_1+bg_2-b\phi)\beta + (c\phi-cg_1+cg_3-c\phi)\gamma \\ & [\text{subtracting } b(\phi-g_2)\beta = 0 \quad \text{and} \quad c(\phi-g_3)\gamma = 0 \\ = &b(g_2-g_1)\beta + c(g_3-g_1)\gamma. \end{split}$$

161. Thus we see that the operator  $(\phi - g_1)$  cuts off from the general vector  $\xi$  the component parallel to  $\alpha$ .

Operating again, with  $(\phi - g_2)$ , this becomes

$$(\phi-g_1)(\phi-g_2)\xi=c(g_3-g_1)(g_3-g_1)\gamma$$
.

In the same way,

$$(\phi - g_1)(\phi - g_3)\xi = b(g_2 - g_1)(g_2 - g_3)\beta,$$
  
 $(\phi - g_2)(\phi - g_3)\xi = a(g_1 - g_2)(g_1 - g_3)\alpha.$ 

That is, the operator,

$$(\phi - g_2)(\phi - g_3),$$

operating on  $\alpha$ , since  $\xi$  is any vector, leaves its direction unchanged, and

$$\alpha \| (\phi - g_2) (\phi - g_3) \xi$$
.

If the three roots  $g_1$ ,  $g_2$ ,  $g_3$  are unequal the three unchanged directions are given by the operators above.

162. Those lines that are unchanged in direction by the strain  $\phi$  have been suggestively called **latent lines** of  $\phi$ , and evidently from the relation  $(\phi - g_1)\rho = 0$ , the *latent lines of*  $\phi$  are the null directions of  $(\phi - g_1)$ .

Hence  $\phi - g_1$  is a null strain, or § 154,

mod. 
$$(\phi - g_1) = m_g = m - m_1 g + m_2 g^2 - g^3 = 0$$
.

The roots of this cubic,  $g_1, g_2, g_3$  are the ratios of dilatation or extension of the latent lines of  $\phi$  and are called the latent roots of  $\phi$ . Planes whose vectors are not strained out of the plane by  $\phi$  are called the latent planes of  $\phi$ .

**163.** If  $\alpha$ ,  $\beta$ ,  $\gamma$  are the latent directions of  $\phi$ , with the latent roots  $g_1$ ,  $g_2$ ,  $g_3$ , then  $(\beta, \gamma)$ ,  $(\gamma, \alpha)$ ,  $(\alpha, \beta)$  determine the latent planes of  $\phi$ , which are enlarged by the strain  $\phi$  in the ratios  $g_2g_3$ ,  $g_3g_1$ ,  $g_1g_2$ .

For if  $x\beta + y\gamma$  be a vector in the plane  $\beta\gamma$ , then

$$\phi(x\beta+y\gamma)=x\phi\beta+y\phi\gamma=xg_2\beta+yg_3\gamma.$$

and evidently the strained vector remains in the plane, though its direction has been changed. Q.E.D.

Also for the two vectors in the plane  $\beta$ ,  $\gamma$ ,

$$TV\phi(x\beta+y\gamma)(x_1\beta+y_1\gamma)=TV(xg_2\beta+yg_3\gamma)(x_1g_2\beta+y_1g_3\gamma)$$

=
$$g_2g_3TV(x\beta+y\gamma)(x_1\beta+y_1\gamma)$$
 Q.E.D. Conf. § 58.]

**164.** If  $g_2 = g_3$ , and we operate upon upon  $\xi$  (§ 160) with  $\phi - g_2$ , remembering that  $(\phi - g_2)\beta = (\phi - g_2)\gamma = 0$ , we get

$$(\phi - g_2)\xi = a(g_1 - g_2)\alpha,$$

and the latent direction is given by

$$(\phi-g_2)\xi$$
.

Operating upon  $\xi$  by  $(\phi - g_1)$  we get

$$(\phi - g_1)\xi = (g_2 - g_1)(b\beta + c\gamma).$$

Since 
$$(\phi - g_2)(b\beta + c\gamma) = (\phi\beta - g_2\beta) + c(\phi - g_2)\gamma$$
,  
=0  $[(\phi - g_2)\beta = 0, (\phi - g_2)\gamma = 0.$ 

therefore every line of the plane  $\beta_1\gamma$  is unchanged in direction, as well as kept in the plane.

**165.**  $g_1 = g_2 = g_3$ .

Operating on  $\xi = a\alpha + b\beta + cg$  with  $\phi - g_1$  we obtain

$$(\phi - g_1)\xi = 0$$
,

that is, when the latent roots are all equal, all the vectors are latent vectors.

166. There cannot be two latent directions for the same root  $g_1$ . For if  $(\phi - g_1)\delta = 0$  as well as  $(\phi - g_1)\alpha = 0$ , we should get by the method of § 161,

$$(\phi-g_2)(\phi-g_3)\xi=d(g_1-g_2)(g_1-g_3)\delta$$

or 
$$a(g_1-g_2)(g_1-g_3)\alpha = d(g_1-g_2)(g_1-g_3)\delta$$

which cannot be unless

$$g_1 = g_2, \quad g_1 = g_3,$$

that is, unless the roots are all equal.

167. Since the latent planes are strained in different ratios (the latent roots being unequal) they cannot coincide and their intersections, the latent vectors,  $\alpha$ ,  $\beta$ ,  $\gamma$ , cannot be coplanar.

 $\phi - g_1$  strains all vectors into the plane  $\beta$ ,  $\gamma$ , that is, into the plane determined by the other latent vectors.

For if 
$$\xi = x\alpha + y\beta + zy =$$
 general vector,  
then  $(\phi - g_1)\xi = x(\phi - g_1)\alpha + y(\phi - g_1)\beta + z(\phi - g_1)\gamma$   
 $= y(g_2 - g_1)\beta + z(g_3 - g_1)\gamma$ ,  $[(\phi - g_1)\alpha = 0$ .  
=vector in plane  $\beta$ ,  $\gamma$ . Q.E.D.

Repeating this operation we get

$$(\phi - g_1)(\phi - g_2)(\phi - g_3)\xi = 0$$
,

as already shown in § 159.

The strain plane of  $\phi - g_1$  is the latent plane of  $\phi$ .

**168.** Two conjugate strains have the same latent roots. For since  $(\phi - q_1)\alpha = 0$ .

$$\therefore 0 = S\xi(\phi - g_1)\alpha = S\alpha(\phi' - g_1)\xi, \quad [\S 135.$$

and evidently  $\phi'-g_1$  strains all vectors into a plane perpendicular to  $\alpha$ , i.e.,  $\phi'-g_1$  is a null function and  $g_1$  is a latent root of  $\phi'$  (§ 162).

169. The latent plane of one strain is perpendicular to the corresponding latent line of the conjugate strain.

The strain plane of  $\phi'-g_1$  is the latent plane of  $\phi'$ , § 167, and therefore the latent plane of  $\phi'$  is perpendicular to  $\alpha$ , the latent line of  $\phi$ .

170. If a strain is self conjugate its three latent roots are real.

Suppose  $g_1 + t_1 \sqrt{-1}$  to be one of the roots, and let the application of the other factors of the cubic be denoted by

$$(\phi-g_2)(\phi-g_3)\xi=\alpha+\alpha_1\sqrt{-1}.$$

Then, by § 160 (1),

$$\phi(\alpha+\alpha_1\sqrt{-1})=(g_1+t_1\sqrt{-1})(\alpha+\alpha_1\sqrt{-1})$$

or equating the reals and imaginaries,

$$\phi \alpha = g_1 \alpha - t_1 \alpha_1, \quad \phi \alpha_1 = g_1 \alpha_1 + t_1 \alpha.$$

Whence

$$S\alpha_1\phi\alpha = g_1S\alpha_1\alpha - t_1S\alpha_1\alpha_1$$

$$S\alpha\phi\alpha_1 = g_1S\alpha\alpha_1 + t_1S\alpha\alpha$$
,

or, since

$$S\alpha_1\phi\alpha = S\alpha\phi\alpha_1$$

$$0=t_1(\alpha^2+\alpha_1^2),$$

whence

$$t_1 = 0$$
.

Q.E.D.

171. If a strain is self conjugate, that is, if  $\phi = \phi'$  or  $S\eta\phi\xi = S\xi\phi\eta$ , then by § 170 it must have three mutually perpendicular latent directions.

Conversely: If  $\phi$  have three mutually perpendicular latent directions i, j, k, with the corresponding latent roots, a, b, c, then  $\phi$  is self conjugate. For, § 122 (78),

$$\begin{split} \xi &= -\left(iSi\xi + jSj\xi + kSk\xi\right), \\ \phi \xi &= -\left(aiSi\xi + bjSj\xi + ckSk\xi\right); & [\phi i = ai, \text{ etc.} \\ Sn\phi \xi &= -aSniSi\xi - bSniSi\xi - cSnkSk\xi; \end{split}$$

$$\begin{split} \eta &= -\left(iSi\eta + jSj\eta + kSk\eta\right);\\ \phi \eta &= -aiSi\eta - bjSj\eta - ckSk\eta;\\ S\xi \phi \eta &= -aSi\xi Si\eta - bSj\xi Sj\eta - cSk\xi Sk\eta\\ &= S\eta \phi \xi, \end{split}$$

and  $\phi$  is self conjugate.

Q.E.D.

172. When the strain is self conjugate  $(\phi = \phi')$  and the latent roots equal, the strain is non-rotational and is called pure.

Since the strain is self conjugate, § 168, the latent lines are i, j, k, and

$$\begin{split} \xi &= -iSi\xi - jSj\xi - kSk\xi, \\ \phi \xi &= -aiSi\xi - ajSj\xi - akSk\xi \\ &= a(-iSi\xi - jSj\xi - kSk\xi) \\ &= a\xi, \end{split}$$
 [a = latent root.

and  $\xi$  is not rotated.

That is, all vectors are latent lines.

Q.E.D.

When  $\phi$  is self conjugate  $(\phi - \phi')$ , by reference to §§ 143, 144, if the strain is given by

$$\phi i = xi + yj + zk;$$
  

$$\phi j = x'i + y'j + z'k;$$
  

$$\phi k = x''i + y''j + z''k;$$

then when the strain is pure,

$$x'=y, \qquad x''=z, \qquad y''=z',$$

and a pure strain depends upon six instead of upon nine scalar constants.

173. If the strain is self conjugate  $(\phi = \phi')$  and the three latent roots a, b, c are not equal, only the vectors i, j, k are non-rotated.

As in the previous section,

$$\xi = -iSi\xi - jSj\xi - kSk\xi,$$

$$\phi \xi = -aiSi\xi - bjSj\xi - ckSk\xi$$

$$\neq x\xi.$$

$$\phi i = -aiSii - bjSij - ckSik$$

$$= ai.$$
Q.E.D.

Similarly,  $\phi j = bj$ ,  $\phi k = ck$ .

Q.E.D.

174. Example. If  $\phi$  have three latent directions  $\alpha$ ,  $\beta$ ,  $\gamma$  with the three latent roots a, b, c, then

$$\xi = \frac{\alpha S \beta \gamma \xi + \beta S \alpha \gamma \xi + \gamma S \beta \alpha \xi}{S \alpha \beta \gamma}.$$

$$\phi \xi = \frac{a \alpha S \beta \gamma \xi + \dots}{S \alpha \beta \gamma}.$$

$$\phi' \xi = \frac{a \alpha S \beta \gamma \xi + \dots}{S \alpha \beta \gamma},$$
[§ 168.

and  $\phi = \phi'$ , or  $\phi$  is self conjugate. Where is the error?

Ans.  $\phi' \alpha \neq a\alpha$ .

175. Examples. Solve  $V\alpha\xi\beta = \gamma = \phi\xi$  for  $\xi$ . By § 146 (2),

$$m = \frac{1}{S \cdot \lambda \mu \nu} S(V \alpha \lambda \beta V \alpha \mu \beta V \alpha \nu \beta),$$

where  $\lambda$ ,  $\mu$ ,  $\nu$  are any three non-coplanar vectors. Putting for these  $\alpha$ ,  $\beta$ ,  $\gamma$ , if they are not coplanar, we get

$$\begin{split} m = & \frac{1}{S \cdot \alpha \beta \gamma} S(\alpha^2 \beta \cdot \alpha \beta^2 V \alpha \gamma \beta) = & \frac{\alpha^2 \beta^2}{S \cdot \alpha \beta \gamma} S(\beta \alpha V \alpha \gamma \beta) = \alpha^2 \beta^2 S \alpha \beta. \\ & [V \alpha \gamma \beta = & \alpha S \gamma \beta - \gamma S \alpha \beta + \beta S \alpha \gamma. \quad [\S 122 (64). \end{split}$$

By § 158,

$$m_1 = \frac{1}{S\alpha\beta\gamma} S(\alpha \cdot \alpha\beta^2 \cdot V\alpha\gamma\beta + \beta V\alpha\gamma\beta \cdot \alpha^2\beta + \gamma \cdot \alpha^2\beta \cdot \alpha\beta^2)$$
$$= -\alpha^2\beta^2.$$

$$m_2 = -S\alpha\beta.$$

By § 157 (1), we get

$$\begin{split} \alpha^2\beta^2S\alpha\beta\cdot\phi^{-1}\gamma &= -\alpha^2\beta^2\gamma + S\alpha\beta V\alpha\gamma\beta + V(\alpha V\alpha\gamma\beta\cdot\beta) \\ &= -\alpha^2\beta^2\gamma + S\alpha\beta V\alpha\gamma\beta + \alpha^2S\gamma\beta\cdot\beta \\ &\quad - S\alpha\beta V\alpha\gamma\beta + S\alpha\gamma\cdot V\alpha\beta^2 \\ &\quad [\S~122~(64). \end{split}$$

$$= -\alpha^2 \beta^2 \gamma + \alpha^2 \beta S \beta \gamma + \alpha \beta^2 S \alpha \gamma,$$

whence

$$\phi^{-1}\gamma = \xi = \frac{-\gamma + \alpha^{-1}S\alpha\gamma + \beta^{-1}S\beta\gamma}{S\alpha\beta}.$$

## APPENDIX

#### FUNCTIONAL SYMBOLS

As every discrete magnitude is of necessity derived from unity by some algebraic operation, so the symbol representing a discrete magnitude can be considered as the symbol of some operation whose operand is unity.

Symbolizing the operation of converting unity into the magnitude x by the symbol  $x \cdot 1$ , we have the functional equation (x operating upon 1),

$$x \cdot 1 = x$$
.

A second application gives

$$x(x\cdot 1)=x^2\cdot 1=x^2,$$

where the 2 of  $x^2 \cdot 1$  shows the number of operations, and the 2 of  $x^2$  the result.

The inverse of the operation must be symbolized by

$$x^{-1}\cdot 1 = \frac{1}{x}\cdot 1.$$

In the expression  $x \cdot 1$  occur two concepts, the operation,  $x \cdot 1$  and the effect of the operation, x.

The inverse of the operation is  $x^{-1} \cdot 1 = \frac{1}{x} \cdot 1$ , the inverse of the effect is  $(x)^{-1} = \frac{1}{x}$ , and in this case the two results coincide.

If we take a different operand, say y, then  $x^{-1} \cdot y$  and  $(xy)^{-1}$ , the inverse of the operation and the inverse of the effect are not the same.

Every algebraic expression can be considered as a functional operation. Thus adding 1 to x can be considered as an operation symbolized thus,

$$f(x) = x + 1.$$

The inverse operation would be whatever operation is necessary to change back from x+1 to x to original operand. Thus

$$f^{-1}(x) = x - 1.$$

Here again  $f^{-1} \neq \frac{1}{f}$ , where, of course,  $f^{-1}$  symbolizes the inverse of the operation and  $(f)^{-1} = \frac{1}{f}$ , the inverse of the effect.

Similarly, 
$$\log^{-1} x \neq (\log x)^{-1}$$
,  
 $\sin^{-1} x \neq (\sin x)^{-1}$ 

Examples. If f(x) = x + 1,  $f^{-1}(x) = x - 1$ ,  $f^{2}(x) = x + 2$ .

If 
$$f(x) = \frac{1}{x^2 + 1}$$
,  $f^{-1}(x) = \sqrt{\frac{1 - x}{x}}$ ,  $f^2(x) = \frac{x^4 + 2x^2 + 1}{x^4 + 2x^2 + 2}$ .

If 
$$f(x) = \frac{1}{x} + x$$
,  $f^{-1}(x) = \frac{x - \sqrt{x^2 - 4}}{2}$ ,  $f^2(x) = \frac{x}{1 + x^2} + \frac{1 + x^2}{x}$ .

If 
$$f(x) = x^2 + 2x + 6$$
,  $f^{-1}(x) = -1 \pm \sqrt{x - 5}$ ,  $f^2(x) = (x^2 + 2x + 6)^2 + 2(x^2 + 2x + 6) + 6$ .

In these examples, omitting the operand, show that

$$ff^{-1} = f^{\circ} = 1$$
,  $f^{2}f^{-1} = f$   $f^{-1}f^{2} = f$ ,  $f^{-1}f = f^{\circ} = 1$ ,  
 $f^{3}f^{-2} = f$ ,  $f^{-3}f = f^{-2}$ .

In the same way prove

$$\sin \sin^{-1} x = x, \qquad \log \log^{-1} x = x,$$

$$\log^{-1} \log x = x, \qquad \sin^{-1} \sin^{2} x = \sin x.$$
If  $f(x) = x^{2} + 3$ ,  $F(x) = 2 - \sqrt{x}$ , then
$$fF(x) = (2 - \sqrt{x})^{2} + 3, \qquad F \cdot f(x) = 2 - \sqrt{x^{2} + 3}.$$
If  $\phi(x) = \frac{2x - 1}{3x - 2} = y$ , then  $\phi(y) = x$ ,  $\phi^{2}(x) = x$ .

The algebraic symbol is distributive over and commutative with its operand, that is, xy+xz+...=x(y+z+...), xy=yx.

Other symbols of operation which like these are distributive over and commutative with the operand will be subject to the same algebraic laws. Hence we can treat these symbols of operation just as we treat algebraic symbols of operation.

Hence we can write § 122 (33), (34), § 132, § 135,

$$Sp + Vp = (S + V)p;$$
  $\phi \xi - \phi' \xi = (\phi - \phi') \xi,$   
 $\phi n \xi + \phi' \xi = (n\phi + \phi') \xi,$   $\phi \xi + gn \xi = (\phi + gn) \xi.$ 

We recall the similar results in Calculus, where, omitting the operand, we have

$$(D^2 - a^2) = (D + a)(D - a) \qquad [D = \frac{d}{dx}]$$

$$(D^2 - D - 2) = (D + 1)(D - 2)$$

$$x^2D^2 = x^2D \cdot D = x^2D\frac{\theta}{x} \qquad [xD = \theta]$$

$$= x^2\frac{xd \cdot \theta - \theta}{x^2}$$

$$= \theta \cdot \theta - \theta = \theta(\theta - 1),$$

where  $\theta$  is commutative with constants but not with either  $\boldsymbol{x}$  or  $\boldsymbol{D}$ .

## INDEX

Addition, 3 Algebra, 31 Amplitude, 7 Angle of a quaternion, 39 Argand diagram, 7 Arithmetic, 30 Axis of quaternion, 21 Calculi, 2 Complex functions, 31 Complex quantity, 7 Conjugate of a quaternion, 33 Conjugate-strain function, 65 Continuum, 1 Coplanar quaternion, 46 Diplanar quaternions, 36 Discreta, 1 Division. 4 Euler's Theorem, 28 Evolution, 5 Functional symbols, 87 Harmonic displacement, 11 Idiograph, 6 Idiograph, space, 10 Involution, 4 Latent lines, 79 Latent planes, 79 Linear homogeneous strain, 67 Linear scalar equation, 62 Mean reversion, 5

Modulus, 7, 71 Multiplication, 4 of vectors, 17 Nonion, 70 Null direction, 72 Null function, 72 Operation A, B, 11Opposite quaternions, 33 Parallel vectors, 17 Plane of quaternion, 21 Plane of a vector, 21 Quadrantal versors, 50 Quaternion, 21, 31 , angle of, 39 " , axis of, 21 .. , plane of, 21 , reciprocal of, 32 Quaternions, equality of, 35 , diplanar, 36 Revector, 12 Reversion, 3 Rotator, 52, 68 Scalar, 9 " part, 27 of the unit part, 27 Self-conjugate strain function, 66 Singly null, 72 Space idiograph, 10 Stroke, 6

# INDEX

Strain, 67	Vector, 11
Strain, linear homogeneous, 67	", addition, etc., 12
Strain function, 64	" unit, 13
" conjugate, 65	", inclined, 18
Strain plane, 72	'', parallel, 17
Subtraction, 4	'', perpendicular, 18
Tensor, 13, 27	" part, 27
of the vector part, 27	" part of the unit part, 27
Unit part, 27	'' , plane of, 21
" vector, 13	Versors, 48, 49
" of the vector part, 27	", quadrantal, 50





## D. VAN NOSTRAND COMPANY 23 MURRAY AND 27 WARREN STREETS NEW YORK

#### SHORT-TITLE CATALOG

OF

### Publications and Importations

OF

#### SCIENTIFIC AND ENGINEERING BOOKS



This list includes the technical publications of the following English publishers:

SCOTT, GREENWOOD & CO. CROSBY LOCKWOOD & SON CONSTABLE & COMPANY, Ltd. TECHNICAL PUBLISHING CO. ELECTRICIAN PRINTING & PUBLISHING CO.,

for whom D. Van Nostrand Company are American agents.

Descriptive Circulars sent on request.

#### SHORT-TITLE CATALOG

OF THE

#### **Publications and Importations**

0F

## D. VAN NOSTRAND COMPANY 23 MURRAY AND 27 WARREN STREETS, N. Y.

Prices marked with an asterisk (\*) are NET All bindings are in cloth unless otherwise noted

A B C Code. (See Clausen-Thue.)		
Abbott, A. V. The Electrical Transmission of Energy8vo,	<b>*\$</b> 5	00
— A Treatise on Fuel. (Science Series No. 9.) 16mo,	Ò	50
Testing Machines. (Science Series No. 74.)16mo,	0	50
Adam, P. Practical Bookbinding. Trans. by T. E. Maw. 12mo,	*2	50
Adams, J. W. Sewers and Drains for Populous Districts8vo,	2	50
Addyman, F. T. Practical X-Ray Work8vo,	*4	00
Ar Code. (See Clausen-Thue.)		
Aikman, C. M. Manures and the Principles of Manuring 8vo,	2	50
Alexander, J. H. Elementary Electrical Engineering12mo,	2	00
- Universal Dictionary of Weights and Measures 8vo,	3	50
Allan, W. Strength of Beams under Transverse Loads.		
(Science Series No. 19.)16mo,	0	50
— Theory of Arches. (Science Series No. 11.)16mo,	. 0	50
Allen, H. Modern Power Gas Producer Practice and Applica-	·	
tions12mo,	*2	50
Gas and Oil Engines8vo,	*4	50
Anderson, F. A. Boiler Feed Water8vo,	*2	50
Anderson, Cap. G. L. Handbook for the Use of Electricians .8vo,	3	00
Anderson, J. W. Prospector's Handbook12mo,	I	50

Andes, L. Vegetable Fats and Oils8vo,	*4	00
—— Animal Fats and Oils. Trans. by C. Salter8vo,	*4	00
Drying Oils, Boiled Oil, and Solid and Liquid Driers8vo,	*5	00
—— Iron Corrosion, Anti-fouling and Anti-corrosive Paints.		
Trans. by C. Salter8vo,	*4	00
Oil Colors and Printers' Ink. Trans. by A. Morris and		
H. Robson8vo,	*2	50
Treatment of Paper for Special Purposes. Trans. by C.		
Salter12mo,	*2	50
Annual Reports on the Progress of Chemistry.		
Vol. I. (1904)8vo,	*2	00
Vol. II. (1905)8vo,	*2	00
Vol. III. (1906)8vo,	*2	00
Vol. IV. (1907)8vo,	*2	00
Vol. V. (1908)8vo,	*2	00
Vol. VI. (1909)8vo,	*2	00
Argand, M. Imaginary Quantities. Translated from the French		
by A. S. Hardy. (Science Series No. 52.)16mo,	0	50
Armstrong, R., and Idell, F. E. Chimneys for Furnaces and		
Steam Boilers. (Science Series No. 1.)16mo,	0	50
Arnold, E. Armature Windings of Direct Current Dynamos.		
Trans. by F. B. DeGress8vo,	*2	00
Ashe, S. W., and Keiley, J. D. Electric Railways. Theoreti-		
cally and Practically Treated. Vol. I. Rolling Stock		
12m0,	*2	50
Ashe, S. W. Electric Railways. Vol. II. Engineering Pre-		-
liminaries and Direct Current Sub-Stations12mo,	*2	50
Electricity: Experimentally and Practically Applied.		
12mo,	*2	00
Atkinson, A. A. Electrical and Magnetic Calculations 8vo,	*1	50
Atkinson, J. J. Friction of Air in Mines. (Science Series		-
No. 14.)16mo,	0	50
Atkinson, J. J., and Williams, E. H., Jr. Gases Met with in		
Coal Mines. (Science Series No. 13.)16mo,	0	50
Atkinson, P. The Elements of Electric Lighting 12mo,	I	50
The Elements of Dynamic Electricity and Magnetism. 12mo,		00
Power Transmitted by Electricity	2	00
Auchincloss, W. S. Link and Valve Motions Simplified 8vo,	*1	50
Ayrton, H. The Electric Arc8vo,		00

Bacon, F. W. Treatise on the Richards Steam-Engine Indica-		
tor12m0,	1	00
Bailes, G. M. Modern Mining Practice. Five Volumes, 8vo, each,		50
Bailey, R. D. The Brewers' Analyst	_	00
Baker, A. L. Quaternions12mo,		25
Baker, Benj. Pressure of Earthwork. (Science Series No. 56.)	_	-3
16mo,		
Baker, I. O. Levelling. (Science Series No. 91.)16mo,	0	50
Baker, M. N. Potable Water. (Science Series)16mo,		50
— Sewerage and Sewage Purification. (Science Series No. 18.)	•	50
16mo,	0	50
·	*1	-
Bale, G. R. Modern Iron Foundry Practice. Two Volumes.		~3
12m0.		
Vol. I. Foundry Equipment, Material Used	*,	50
Vol. II. Machine Moulding and Moulding Machines		50
Bale, M. P. Pumps and Pumping12mo,		50
Ball, R. S. Popular Guide to the Heavens		50
— Natural Sources of Power. (Westminster Series)8vo,	*2	
Ball, W. V. Law Affecting Engineers8vo,	*3	
Bankson, Lloyd. Slide Valve Diagrams. (Science Series No.	3	30
108.)	^	50
Barba, J. Use of Steel for Constructive Purposes12mo,		00
Barker, A. Textiles and Their Manufacture. (Wetsminster	. •	00
Series)	•	00
Barker, A. H. Grapic Methods of Engine Design12mo,	*1	
Barnard, F. A. P. Report on Machinery and Processes of the	•	30
Industrial Arts and Apparatus of the Exact Sciences at		
the Paris Universal Exposition, 18678vo,	_	00
Barnard, J. H. The Naval Miliatiaman's Guide16mo, leather,	-	25
Barnard, Major J. G. Rotary Motion. (Science Series No. 90.)	•	<b>~</b> 3
16mo,	^	50
Barrus, G. H. Boiler Tests8vo,	*3	-
Engine Tests8vo,	*4	
The above two purchased together	*6	
Barwise, S. The Purification of Sewage	3	
Baterden, J. R. Timber. (Westmenster Series)8vo,	*2	-
Bates, E. L., and Charlesworth, F. Practical Mathematics and	-	30
Geometry for Technical Students	*1	E۸
Geometry for Accumical Dissembs#	-	30

Beadle, C. Chapters on Papermaking. Five Volumes	
12mo, each,	*2 00
Beaumont, R. Color in Woven Design8vo,	7 50
— Finishing of Textile Fabrics8vo,	*4 00
Beaumont, W. W. The Steam-Engine Indicator8vo,	2 50
Bedell, F., and Pierce, C. A. Direct and Alternating Current	
Testing8vo,	*2 00
Beech, F. Dyeing of Cotton Fabrics8vo,	*3 00
—— Dyeing of Woolen Fabrics8vo,	*3 50
Beckwith, A. Pottery8vo, paper,	o 60
Begtrup, J. The Slide Valve8vo,	*2, 00
Bender, C. E. Continuous Bridges. (Science Series No. 26.)	
16mo,	0 . 50
Proportions of Piers used in Bridges. (Science Series No. 4.)	
16mo,	0 50
Bennett, H.G. The Manufacture of Leather	*4 50
Bernthsen, A. A Text-book of Organic Chemistry. Trans. by	_
G. M'Gowan12mo,	*2 50
Berry, W. J. Differential Equations of the First Species.	
12mo (In Prepar	ration.)
r2mo (In Prepar Bersch, J. Manufacture of Mineral and Lake Pigments. Trans.	ŕ
Bersch, J. Manufacture of Mineral and Lake Pigments. Trans. by A. C. Wright	*5 00
Bersch, J. Manufacture of Mineral and Lake Pigments. Trans. by A. C. Wright	*5 00 5 00
Bersch, J. Manufacture of Mineral and Lake Pigments. Trans. by A. C. Wright	*5 00 5 00 *4 00
Bersch, J. Manufacture of Mineral and Lake Pigments. Trans. by A. C. Wright	*5 00 5 00 *4 00 *5 00
12mo (In Preparation of Mineral and Lake Pigments. Trans. by A. C. Wright	*5 00 5 00 *4 00 *5 00 *7 50
12mo (In Preparation   12mo   12	*5 00 5 00 *4 00 *5 00 *7 50 *2 00
12mo (In Preparation   12mo   12	*5 00 5 00 *4 00 *5 00 *7 50 *2 00 *1 25
12mo (In Preparations)   12mo (In Preparatio	*5 00 5 00 *4 00 *5 00 *7 50 *2 00 *1 25 *1 50
12mo (In Preparation   12mo   12	*5 00 5 00 *4 00 *5 00 *7 50 *2 00 *1 25 *1 50 *4 00
12mo (In Preparation   12mo   12	*5 00 5 00 *4 00 *5 00 *7 50 *2 00 *1 25 *1 50 *4 00 2 00
12mo (In Preparation   12mo   12	*5 00 5 00 *4 00 *5 00 *7 50 *2 00 *1 25 *1 50 *4 00 2 00 *6 00
Bersch, J. Manufacture of Mineral and Lake Pigments. Trans. by A. C. Wright	*5 00 5 00 *4 00 *5 00 *7 50 *2 00 *1 25 *1 50 *4 00 2 00 *6 00 7 50
Bersch, J. Manufacture of Mineral and Lake Pigments. Trans. by A. C. Wright	*5 00 5 00 *4 00 *5 00 *7 50 *2 00 *1 25 *1 50 *4 00 2 00 *6 00 7 50 7 50
Bersch, J. Manufacture of Mineral and Lake Pigments. Trans. by A. C. Wright	*5 00 5 00 *4 00 *5 00 *7 50 *2 00 *1 25 *1 50 *4 00 2 00 *6 00 7 50 *2 50
Bersch, J. Manufacture of Mineral and Lake Pigments. Trans. by A. C. Wright	*5 00 5 00 *4 00 *5 00 *7 50 *2 00 *1 25 *1 50 *4 00 2 00 *6 00 7 50 *2 50 5 00
Bersch, J. Manufacture of Mineral and Lake Pigments. Trans. by A. C. Wright	*5 00 5 00 *4 00 *5 00 *7 50 *2 00 *1 25 *1 50 *4 00 2 00 *6 00 7 50 *2 50 5 00 5 00
Bersch, J. Manufacture of Mineral and Lake Pigments. Trans. by A. C. Wright	*5 00 5 00 *4 00 *5 00 *7 50 *2 00 *1 25 *1 50 *4 00 2 00 *6 00 7 50 *2 50 5 00

Booth, W. H. Superheaters and Superheating and their Con-		
trol8vo,	*I	50
Bottcher, A. Cranes: Their Construction, Mechanical Equip-		_
ment and Working. Trans. by A. Tolhausen 4to,	*10	00
Bottler, M. Modern Bleaching Agents. Trans. by C. Salter		
12m0,	*2	50
Bottone, S. R. Magnetos for Automobilists12mo,		00
Boulton, S. B. Preservation of Timber. (Science Series No.		
82.)r6mo,	0	50
Bourgougnon, A. Physical Problems. (Science Series No. 113.)		•
16mo,	0	50
Bourry, E. Treatise on Ceramic Industries. Trans. by		_
W. P. Rix8vo (In	Pre	88.)
Bow, R. H. A Treatise on Bracing8vo,		50
Bowie, A. J., Jr. A Practical Treatise on Hydraulic Mining .8vo,		00
Bowker, W. R. Dynamo Motor and Switchboard Circuits 8vo,		50
Bowles, O. Tables of Common Rocks. (Science Series) .16mo,		50
Bowser, E. A. Elementary Treatise on Analytic Geometry. 12mo,		75
- Elementary Treatise on the Differential and Integral	_	
Calculus12mo,	2	25
— Elementary Treatise on Analytic Mechanics 12mo,		00
Elementary Treatise on Hydro-mechanics12mo,		50
— A Treatise on Roofs and Bridges		25
Boycott, G. W. M. Compressed Air Work and Diving8vo,		00
Bragg, E. M. Marine Engine Design12mo,	•	00
Brainard, F. R. The Sextant. (Science Series No. 101.). 16mo,		
Brassey's Naval Annual for 19108vo,	*6	00
Brew, W. Three-Phase Transmission8vo,		00
Brewer, R. W. A. The Motor Car12mo,	*2	00
Briggs, R., and Wolff, A. R. Steam-Heating. (Science Series		
No. 67.)16mo,	0	50
Bright, C. The Life Story of Sir Charles Tilson Bright 8vo,		50
British Standard Sections8x15	•	00
Complete list of this series (45 parts) sent on application.		
Broughton, H. H. Electric Cranes and Hoists(In	Pres	88.)
Brown, G. Healthy Foundations. (Science Series No. 80.)		.,
16mo.	0	50
Brown, H. Irrigation8vo,	*5	00
Drawn Wm N The Art of Framelling on Metal	*-	

•		
Brown, Wm. N. Handbook on Japanning and Enamelling. 12mo,	*1	50
House Decorating and Painting12mo,	*1	50
History of Decorative Art12mo,		25
- Dipping, Burnishing, Lacquering and Bronzing Brass		•
Ware	*1	00
— Workshop Wrinkles8vo,	*1	00
Browne, R. E. Water Meters. (Science Series No. 81.). 16mo,	0	50
Bruce, E. M. Pure Food Tests	*1	25
Bruhns, Dr. New Manual of Logarithms8vo, half mor.,	2	50
Brunner, R. Manufacture of Lubricants, Shoe Polishes and		
Leather Dressings. Trans. by C. Salter8vo,	*3	00
Buel, R. H. Safety Valves. (Science Series No. 21.)61mo,	0	50
Bulmann, H. F., and Redmayne, R. S. A. Colliery Working and		
Management8vo,	6	00
Burgh, N. P. Modern Marine Engineering 4to, half mor.,	IO	00
Burt, W. A. Key to the Solar Compass16mo, leather,	2	50
Burton, F. G. Engineering Estimates and Cost Accounts.		
12m0,		50
Buskett, E. W. Fire Assaying12mo,	*1	25
0 · W D140 · 1 · 1 · 0 · 1		
Cain, W. Brief Course in the Calculus12mo,		75
Elastic Arches. (Science Series No. 48.)		50
Maximum Stresses. (Science Series No. 38.)16mo,	0	50
—— Practical Designing Retaining of Walls. (Science Series		
No. 3.)16mo,	0	50
— Theory of Steel-concrete Arches and of Vaulted Structures.		
(Science Series)16mo,		50
— Theory of Voussoir Arches. (Science Series No. 12.) 16mo,		50
Symbolic Algebra. (Science Series No. 73.)16mo,		50
Campin, F. The Construction of Iron Roofs	2	00
Carpenter, F. D. Geographical Surveying. (Science Series		
No. 37.)		
Carpenter, R. C., and Diederichs, H. Internal-Combustion	•	
Engines	*5	00
Carter, E. T. Motive Power and Gearing for Electrical Machin-	*	
ery	*5	
Carter, H. A. Ramie (Rhea), China Grass		00
Carter, H. R. Modern Flax, Hemp, and Jute Spinning 8vo,	*3 *2	

Cathcart, W. L., and Chaffee, J. I. Elements of Graphic Statics.  8vo.	*-	00
•	3	OU
Caven, R. M., and Lander, G. D. Systematic Inorganic Chemis-	_	
try12mo,		00
Chambers' Mathematical Tables8vo,	I	75
Charnock, G. F. Workshop Practice. (Westminster Series.)		
. 8vo (In	Pre	88.)
Charpentier, P. Timber8vo,	*6	00
Chatley, H. Principles and Designs of Aeroplanes. (Science		
Series.)16mo,	0	50
How to Use Water Power 12mo,		00
Child, C. T. The How and Why of Electricity12mo,		00
Christie, W. W. Boiler-waters, Scale, Corrosion, Foaming.8vo,		00
— Chimney Design and Theory	_	00
	_	
Furnace Draft. (Science Series)		50
Church's Laboratory Guide. Rewritten by Edward Kinch8vo,		50
Clapperton, G. Practical Papermaking8vo,		50
Clark, C. H. Marine Gas Engines(In	Pre	8 <b>8.</b> )
Clark, D. K. Rules, Tables and Data for Mechanical Engineers		
8vo,	5	00
Fuel: Its Combustion and Economy12mo,	I	50
The Mechanical Engineer's Pocketbook	2	00
Tramways: Their Construction and Working8vo,		50
Clark, J. M. New System of Laying Out Railway Turnouts	•	-
12m0,	т	00
Clausen-Thue, W. A B C Telegraphic Code. Fourth Edition	-	•
12mo,	*5	~~
Fifth Edition8vo,	_	
	•	00
— The Ar Telegraphic Code	*7	
Cleemann, T. M. The Railroad Engineer's Practice12mo,	*1	50
Clerk, D., and Idell, F. E. Theory of the Gas Engine. (Science		
Series No. 62.)	0	50
Clevenger, S. R. Treatise on the Method of Government Sur-		
veying	2	50
Clouth, F. Rubber, Gutta-Percha, and Balata8vo,	*5	00
Coffin, J. H. C. Navigation and Nautical Astronomy12mo,	*3	50
Colburn, Z., and Thurston, R. H. Steam Boiler Explosions.	_	-
(Science Series No. 2.)16mo,	O	50
Cole P S Treetise on Photographic Ontics Tamo		50

#### 10 d. van nostrand company's short-title catalog

Deerr, N. Sugar and the Sugar Cane8vo,	*3	00
Deite, C. Manual of Soapmaking. Trans. by S. T. King 4to,	*5	00
De la Coux, H. The Industrial Uses of Water. Trans. by A.		
Morris8vo,	*4	50
Del Mar, W. A. Electric Power Conductors8vo,	*2	00
Denny, G. A. Deep-Level Mines of the Rand4to,	*10	00
—— Diamond Drilling for Gold	*5	00
De Roos, J. D. C. Linkages. (Science Series No. 47.)r6mo,	0	50
De Varona, A. Sewer Gases. (Science Series No. 55.) 16mo,		50
Derr, W. L. Block Signal OperationOblong 12mo,	*1	50
Desaint, A. Three Hundred Shades and How to Mix Them 8vo,	*10	00
Dibdin, W. J. Public Lighting by Gas and Electricity8vo,	*8	00
Purification of Sewage and Water8vo,	6	50
Dietrich, K. Analysis of Resins, Balsams, and Gum Resins .8vo,	*3	00
Dinger, Lieut. H. C. Care and Operation of Naval Machinery		
. 12mo.	*2	00
Dixon, D. B. Machinist's and Steam Engineer's Practical Cal-		
culator16mo, mor.,	1	25
Doble, W. A. Power Plant Construction on the Pacific Coast.		
``	Pre	88.)
Dodd, G. Dictionary of Manufactures, Mining, Machinery, and		
the Industrial Arts12mo,	I	50
Dorr, B. F. The Surveyor's Guide and Pocket Table-book.		
16mo, mor.,	2	00
Down, P. B. Handy Copper Wire Table16mo,	*1	00
Draper, C. H. Elementary Text-book of Light, Heat and		
Sound12mo,	I	00
Heat and the Principles of Thermo-dynamics12mo,	I	50
Duckwall, E. W. Canning and Preserving of Food Products.8vo,	*5	00
Dumesny, P., and Noyer, J. Wood Products, Distillates, and		
Extracts8vo,	*4	50
Duncan, W. G., and Penman, D. The Electrical Equipment of		
Collieries 8vo,	*4	00
Duthie, A. L. Decorative Glass Processes. (Westminster		
Series)8vo,	*2	00
Dyson, S. S. Practical Testing of Raw Materials8vo,	*5	00
Eccles, R. G., and Duckwall, E. W. Food Preservatives 8vo,	1	00
paper,	o	50

Eddy, H. T. Researches in Graphical Statics8vo, I	50
Maximum Stresses under Concentrated Loads8vo, I	50
Edgcumbe, K. Industrial Electrical Measuring Instruments.	-
8vo, *2	50
· · · · · · · · · · · · · · · · · · ·	50
and the state of t	50
	00
	00
	00
	00
Ekin, T. C. Water Pipe and Sewage Discharge Diagrams. folio, *3	
Eliot, C. W., and Storer, F. H. Compendious Manual of Qualita-	00
	-
	00
Ennis, Wm. D. Linseed Oil and Other Seed Oils8vo, *4	
— Applied Thermodynamics8vo, *4	
Flying Machines To-day	•
Erfurt, J. Dyeing of Paper Pulp. Trans. by J. Hubner8vo, *7	_
Erskine-Murray, J. A Handbook of Wireless Telegraphy 8vo, *3	
Evans, C. A. Macadamized Roads	
Ewing, A. J. Magnetic Induction in Iron8vo, *4	00
Fairie, J. Notes on Lead Ores 12mo, *1	
Notes on Pottery Clays12mo, *1	50
Fairley, W., and Andre, Geo. J. Ventilation of Coal Mines.	
	50
Fairweather, W. C. Foreign and Colonial Patent Laws8vo, *3	00
Fanning, T. T. Hydraulic and Water-supply Engineering.	
8vo, *5	00
Fauth, P. The Moon in Modern Astronomy. Trans. by J.	
McCabe8vo, *2	00
Fay, I. W. The Coal-tar Colors	s.)
Fernbach, R. L. Glue and Gelatine8vo, *3	00
Fischer, E. The Preparation of Organic Compounds. Trans.	
by R. V. Stanford 12mo, *1	25
ti t	00
Fisher, H. K. C., and Darby, W. C. Submarine Cable Testing.	
8vo, *3	50
, ,	50
	_

Fleischmann, W. The Book of the Dairy. Trans. by C. M. Aikman		. 00
Fleming, J. A. The Alternate-current Transformer. Two	4	
Volumes8vo,		
Vol. I. The Induction of Electric Currents	_	00
Vol. II. The Utilization of Induced Currents		00
—— Centenary of the Electrical Current8vo,	*0	50
Electric Lamps and Electric Lighting8vo,	*3	00
— Electric Laboratory Notes and Forms4to,	*5	00
A Handbook for the Electrical Laboratory and Testing		
Room. Two Volumes8vo, each,	*5	00
Fluery, H. The Calculus Without Limits or Infinitesimals.	•	
Trans. by C. O. Mailloux(In	Pre	88.)
Flynn, P. J. Flow of Water. (Science Series No. 84.) 16mo,		50 50
— Hydraulic Tables. (Science Series No. 66.) 16mo,		50
	U	50
Foley, N. British and American Customary and Metric Meas-	<b>.</b>	
uresfolio,	-3	00
Foster, H. A. Electrical Engineers' Pocket-book. (Sixth		
Edition.) 12mo, leather,	5	00
Foster, Gen. J. G. Submarine Blasting in Boston (Mass.)		
Harbor4to,	3	50
Fowle, F. F. Overhead Transmission Line Crossings 12mo,		50
The Solution of Alternating Current Problems.		•
8vo (In	Pres	(_R)
Fox, W. G. Transition Curves. (Science Series No. 110.).16mo,		50
Fox, W., and Thomas, C. W. Practical Course in Mechanical	·	30
Drawing12mo,	_	
		25
Foye, J. C. Chemical Problems. (Science Series No. 69.).16mo,		50
— Handbook of Mineralogy. (Science Series No. 86.). 16mo,		50
Francis, J. B. Lowell Hydraulic Experiments4to,	15	
Frye, A. I. Civil Engineers' Pocket-book(In	Pres	8.)
Fuller, G. W. Investigations into the Purification of the Ohio		
River4to,		
Furnell, J. Paints, Colors, Oils, and Varnishes8vo,	*1	00
Gant, L. W. Elements of Electric Traction8vo,	*2	50
Garcke, E., and Fells, J. M. Factory Accounts8vo,	3	_
Garforth, W. E. Rules for Recovering Coal Mines after Explo-	•	
gions and Fires		EO.

Gaudard, J. Foundations. (Science Series No. 34.)16mo, Gear, H. B., and Williams, P. F. Electric Central Station Dis-		50
tributing Systems	ratio	n.)
Geerligs, H. C. P. Cane Sugar and Its Manufacture8vo,		00
Geikie, J. Structural and Field Geology8vo,		00
Gerber, N. Analysis of Milk, Condensed Milk, and Infants'	7	•
Milk-Food		25
Gerhard, W. P. Sanitation, Water-supply and Sewage Disposal	•	43
of Country Houses12mo,	*-	00
Gas Lighting. (Science Series No. 111.)16me		50
—— Household Wastes. (Science Series No. 97.) 16mo.		~
House Drainage. (Science Series No. 63.)16mo,		50
	0	50
Sanitary Drainage of Buildings. (Science Series No. 93.)		
16mo,		50
Gerhardi, C. W. H. Electricity Meters8vo,	*4	00
Geschwind, L. Manufacture of Alum and Sulphates. Trans.		
by C. Salter8vo,	*5	00
Gibbs, W. E. Lighting by Acetylene12mo,	*1	50
Physics of Solids and Fluids. (Carnegie Technical Schools		
Text-books.)	*x'	50
Gibson, A. H. Hydraulics and Its Application8vo,	*5	00
Water Hammer in Hydraulic Pipe Lines12mo,	*2	00
Gilbreth, F. B. Motion Study. A Method for Increasing the		
Efficiency of the Workman12mo,	*2	00
Gillmore, Gen. Q. A. Limes, Hydraulic Cements and Mortars.	_	•
8vo.	4	00
Roads, Streets, and Pavements12mo,	•	00
Golding, H. A. The Theta-Phi Diagram12mo,		25
Goldschmidt, R. Alternating Current Commutator Motor .8vo.		43 00
Goodchild, W. Precious Stones. (Westminster Series.)8vo,		
		00
Goodeve, T. M. Textbook on the Steam-engine12mo,		00
Gore, G. Electrolytic Separation of Metals8vo,		50
Gould, E. S. Arithmetic of the Steam-engine12mo,		00
— Calculus. (Science Series No. 112.)		50
— High Masonry Dams. (Science Series No. 22.)16mo,	0	50
—— Practical Hydrostatics and Hydrostatic Formulas. (Science		
Series.)16mo,		50
Grant, J. Brewing and Distilling. (Westminster Series.) 8vo (In	Pres	ss.)
Grav. J. Electrical Influence Machines	2	00

Greenwood, E. Classified Guide to Technical and Commercial	_	
Books8vo,	_	00
Gregorius, R. Mineral Waxes. Trans. by C. Salter 12mo,	_	00
Griffiths, A. B. A Treatise on Manures12mo,	3	00
— Dental Metallurgy8vo,	*3	50
Gross, E. Hops8vo,	*4	50
Grossman, J. Ammonia and its Compounds12mo,	*1	25
Groth, L. A. Welding and Cutting Metals by Gases or Electric-		
ity8vo,	**3	00
Grover, F. Modern Gas and Oil Engines8vo,	*2	00
Gruner, A. Power-loom Weaving8vo,	*3	00
Güldner, Hugo. Internal-Combustion Engines. Trans. by	•	
	*10	00
Gunther, C. O. Integration12mo,		25
Gurden, R. L. Traverse Tables folio, half mor.		50
Guy, A. E. Experiments on the Flexure of Beams8vo,		25
	_	-3
Haeder, H. Handbook on the Steam-engine. Trans. by H. H.		
P. Powles	3	00
Hainbach, R. Pottery Decoration. Trans. by C. Slater. 12mo,	_	00
Hale, W. J. Calculations of General Chemistry12mo,	•	00
Hall, C. H. Chemistry of Paints and Paint Vehicles12mo,		00
Hall, R. H. Governors and Governing Mechanism 12mo,		00
Hall, W. S. Elements of the Differential and Integral Calculus	_	•
8vo.	*2	25
—— Descriptive Geometry 8vo volume and 4to atlas,	*3	_
Haller, G. F., and Cunningham, E. T. The Tesla Coil 12mo,	*1	-
		50
Halsey, F. A. Slide Valve Gears		-
The Use of the Slide Rule. (Science Series.)16mo,		50
Worm and Spiral Gearing. (Science Series.)16mo,		50
Hamilton, W. G. Useful Information for Railway Men. 16mo,		00
Hammer, W. J. Radium and Other Radioactive Substances, 8vo,	*1	
Hancock, H. Textbook of Mechanics and Hydrostatics8vo,		50
Hardy, E. Elementary Principles of Graphic Statics 12mo,	*1	•
Harper, W. B. Utilization of Wood Waste by Distillation. 4to,	*3	
Harrison, W. B. The Mechanics' Tool-book 12mo,		50
Hart, J. W. External Plumbing Work8vo,	*3	
— Hints to Plumbers on Joint Wiping 8vo,	*3	
Principles of Hot Water Supply8vo,	*3	00

Hart, J. W. Sanitary Plumbing and Drainage8vo,	*3	00
Haskins, C. H. The Galvanometer and Its Uses16mo,	1	50
Haskins, C. H. The Galvanometer and Its Uses16mo, Hatt, J. A. H. The Coloristsquare 12mo,		50
Hausbrand, E. Drying by Means of Air and Steam. Trans.		
by A. C. Wright 12mo,	*2	00
Evaporing, Condensing and Cooling Apparatus. Trans.		
by A. C. Wright8vo,	*5	00
Hausner, A. Manufacture of Preserved Foods and Sweetmeats.	_	
Trans. by A. Morris and H. Robson 8vo,	*3	00
Hawke, W. H. Premier Cipher Telegraphic Code4to,	*5	00
100,000 Words Supplement to the Premier Code4to,	_	00
Hawkesworth, J. Graphical Handbook for Réniforced Concrete	·	
Design	*2	50
Hay, A. Alternating Currents8vo,		50
— Principles of Alternate-current Working12mo,		90
— Electrical Distributing Networks and Distributing Lines.8vo,		50
Continuous Current Engineering	_	50
Heap, Major D. P. Electrical Appliances		00
Heaviside, O. Electromagnetic Theory. Two volumes.	-	
8vo. each.	*5	00
Heck, R. C. H. Steam-Engine and Other Steam Motors. Two	3	•
Volumes.		
Vol. I. Thermodynamics and the Mechanics8vo,	*-	50
Vol. II. Form, Construction and Working8vo,	-	90
Abridged edition of above volumes (Elementary)	3	00
8vo (In Prepa		۱,۵
		•
Notes on Elementary Kinematics8vo, boards,		00
Graphics of Machine Forces		00
Hedges, K. Modern Lightning Conductors	3	00
Heermann, P. Dyers' Materials. Trans. by A. C. Wright.		
12m0,	₹2	50
Hellot, Macquer and D'Apligny. Art of Dyeing Wool, Silk and		
Cotton8vo,		00
Henrici, O. Skeleton Structures8vo,		50
Hermann, F. Painting on Glass and Porcelain8vo,	<b>∓</b> 3	50
Hermann, G. The Graphical Statics of Mechanism. Trans.		
by A. P. Smith 12mo,		00
Herzfeld, J. Testing of Yarns and Textile Fabrics8vo,		50
Hildebrandt A. Airshing Past and Present	*3	50

Hildenbrand, B. W. Cable-Making. (Science Series No., 32.)		
16то,		
Hildrich, H., Short History of Chemistry(In	Pre	88.)
Hill, J. W. The Purification of Public Water Supplies. New		
Edition(In		
Interpretation of Water Analysis(In	Pre	ss.)
Hiroi, I. Plate Girder Construction. (Science Series No. 95.)		
16mo,	0	50
Statically-Indeterminate Stresses	*2	00
Hirshfeld, C. F. Engineering Thermodynamics. (Science		
Series.)16mo,	0	50
Hobart, H. M. Heavy Electrical Engineering8vo,		50
Electricity	*2	00
—— Electric Trains8vo,	*2	50
Hobbs, W. R. P. The Arithmetic of Electrical Measurements		
	0	50
Hoff, J. N. Paint and Varnish Facts and Formulas12mo,	*1	50
Hoff, Com.W. B. The Avoidance of Collisions at Sea. 16mo, mor.,	0	<b>7</b> 5
Hole, W. The Distribution of Gas8vo,	*7	50
Holley, A. L. Railway Practicefolio,	12	00
Holmes, A. B. The Electric Light Popularly Explained.		
12mo, paper,		50
Hopkins, N. M. Experimental Electrochemistry8vo,	_	00
Model Engines and Small Boats	I	25
Hopkinson, J., Shoolbred, J. N., and Day, R. E. Dynamic		
Electricity. (Science Series No. 71.)16mo,	. 0	_
Horner, J. Engineers' Turning8vo,	*3	
— Metal Turning12mo,		50
— Toothed Gearing		25
Houghton, C. E. The Elements of Mechanics of Materials. 12mo,	*2	
Houllevigue, L. The Evolution of the Sciences8vo,		
Howe, G. Mathematics for the Practical Man12mo,	*1	25
Howorth, J. Repairing and Riveting Glass, China and Earthen-	_	
ware8vo, paper,	*0	-
Hubbard, E. The Utilization of Wood-waste8vo,	*2	
Humber, W. Calculation of Strains in Girders12mo,	.3	50
Humphreys, A. C. The Business Features of Engineering	*-	
Practice	*I	•

Hurst, G.H. Dictionary of Chemicals and Raw Products. 8vo,	*3	00
— Lubricating Oils, Fats and Greases 8vo,	*3	00
Soaps8vo,	*5	00
—— Textile Soaps and Oils8vo,	*2	50
Hurst, H. E., and Lattey, R. T. Text-book of Physics8vo,	*3	00
Hutchinson, R. W., Jr. Long Distance Electric Power Trans-	_	
mission 12mo,	*3	00
Hutchinson, R. W., Jr., and Ihlseng, M. C. Electricity in	•	
Mining12mo (In	Pres	ss.)
Hutchinson, W. B. Patents and How to Make Money Out of		,
Them12mo,	1	25
Hutton, W. S. Steam-boiler Construction8vo,		00
Practical Engineer's Handbook 8vo,		00
The Works' Manager's Handbook8vo,	-	00
Hyde, E. W. Skew Arches. (Science Series No. 15.)16mo,		50
		•
Induction Coils. (Science Series No. 53.)	0	50
Ingle, H. Manual of Agricultural Chemistry8vo,		00
Innes, C. H. Problems in Machine Design	_	00
— Air Compressors and Blowing Engines	*2	
— Centrifugal Pumps		00
— The Fan		00
Isherwood, B. F. Engineering Precedents for Steam Machinery	_	••
8vo.	2	50
Ivatts, E. B. Railway Management at Stations8vo,		50
Trucing 21 21 Zunitruy annungument at buttons	-	30
Jacob, A., and Gould, E. S. On the Designing and Construction		
of Storage Reservoirs. (Science Series No. 6.). 16mo,	0	50
Iamieson, A. Text Book on Steam and Steam Engines. 8vo,		00
— Elementary Manual on Steam and the Steam Engine . 12mo,	-	50
Jannettaz, E. Guide to the Determination of Rocks. Trans.	•	50
by G. W. Plympton		50
Jehl, F. Manufacture of Carbons	*4	_
Jennings, A. S. Commercial Paints and Painting. (West-	4	00
minster Series.)	Pro	٠. ١
Jennison, F. H. The Manufacture of Lake Pigments 8vo,		
Jepson, G. Cams and the Principles of their Construction8vo,		
• • •		•
Mechanical Drawing		
		w

Johnson, G. L. Photographic Optics and Color Photography	•-	
8vo,	+3	00
Johnson, W. H. The Cultivation and Preparation of Para		
Rubber 8vo,	*3	00
Johnson, W. McA. The Metallurgy of Nickel(In Prepo	ırati	m.)
Johnston, J. F. W., and Cameron, C. Elements of Agricultural		
Chemistry and Geology		60
Joly, J. Radioactivity and Geology12mo,	*3	00
Jones, H. C. Electrical Nature of Matter and Radioactivity		
12m0,	2	00
Jones, M. W. Testing Raw Materials Used in Paint12mo,	*2	00
Jones, L., and Scard, F. I. Manufacture of Cane Sugar8vo,	*5	00
Joynson, F. H. Designing and Construction of Machine Gear-		
ing8vo,	2	00
Jüptner, H. F. V. Siderology: The Science of Iron 8vo,	*5	00
Kansas City Bridge4to,	6	00
Kapp, G. Alternate Current Machinery. (Science Series No.	-	
96.)16mo,	0	50
Dynamos, Motors, Alternators and Rotary Converters.		0-
Trans. by H. H. Simmons8vo.	4	00
— Electric Transmission of Energy	•	50
Keim, A. W. Prevention of Dampness in Buildings 8vo,		00
Keller, S. S. Mathematics for Engineering Students.	_	-
12mo, half leather,		
—— Algebra and Trigonometry, with a Chapter on Vectors	*1	
—— Special Algebra Edition	*1	
• •	*1	
—— Plane and Solid Geometry		_
Analytical Geometry and Calculus	*2	00
Kelsey, W. R. Continuous-current Dynamos and Motors.	_	
8vo,	*2	50
Kemble, W. T., and Underhill, C. R. The Periodic Law and the	_	
Hydrogen Spectrum8vo, paper,	*0	_
Kemp, J. F. Handbook of Rocks8vo,	*1	•
Kendall, E. Twelve Figure Cipher Code4to,	*15	00
Kennedy, A. B. W., and Thurston, R. H. Kinematics of		
Machinery. (Science Series No. 54.)16mo,	0	50
Kennedy, A. B. W., Unwin, W. C., and Idell, F. E. Compressed		
Air. (Science Series No. 106.)16mo,	0	50

Kennedy, R. Modern Engines and Power Generators. Six	
Volumes4to,	15 00
Single Volumes each,	3 00
Electrical Installations. Five Volumes 4to,	15 00
Single Volumes each,	3 50
- Flying Machines; Practice and Design12mo,	*2 00
Kennelly, A. E. Electro-dynamic Machinery8vo,	I 50
Kent, W. Strength of Materials. (Science Series No. 41.).16mo,	0 50
Kershaw, J. B. C. Fuel, Water and Gas Analysis8vo,	*2 50
— Electrometallurgy. (Westminster Series.)8vo,	*2 00
— The Electric Furnace in Iron and Steel Production12mo,	*I 50
Kingdon, J. A. Applied Magnetism8vo,	*3 00
Kinzbrunner, C. Alternate Current Windings8vo,	*1 50
— Continuous Current Armatures8vo,	*I 50
Testing of Alternating Current Machines	*2 00
Kirkaldy, W. G. David Kirkaldy's System of Mechanical	
Testing4to,	10 00
Kirkbride, J. Engraving for Illustration8vo,	*1 50
Kirkwood, J. P. Filtration of River Waters4to,	7 50
Klein, J. F. Design of a High speed Steam-engine8vo,	*5 00
Physical Significance of Entropy8vo,	*1 50
Kleinhans, F. B. Boiler Construction8vo,	3 00
Knight, Capt. A. M. Modern Steamship 8vo,	*7 50
Half Mor.	*9 00
Knox, W. F. Logarithm Tables (In Prepar	ration.)
Knott, C. G., and Mackay, J. S. Practical Mathematics 8vo,	2 00
Koester, F. Steam-Electric Power Plants4to,	*5 00
	*5 00
Koller, T. The Utilization of Waste Products8vo,	*3 50
—— Cosmetics	*2 50
Krauch, C. Testing of Chemical Reagents. Trans. by J. A.	_ 50
Williamson and L. W. Dupre	*3 00
·	3 00
Lambert, T. Lead and its Compounds8vo,	*3 50
— Bone Products and Manures	*3 00
Lamborn, L. L. Cottonseed Products8vo,	*3 00
— Modern Soaps, Candles, and Glycerin8vo,	*7 50
Lamprecht, R. Recovery Work After Pit Fires. Trans. by	, ,,,
C. Salter	*4 00

Lanchester, F. W. Aerial Flight. Two Volumes. 8vo.		
Vol. I. Aerodynamics	*6	0
Vol. II. Aerodonetics	*6	0
Larner, E. T. Principles of Alternating Currents12mo,	*1	2
Larrabee, C. S. Cipher and Secret Letter and Telegraphic	_	
Codei6mo,	0	6
La Rue, B. F. Swing Bridges. (Science Series No. 107.) 16mo,		59
Lassar-Cohn, Dr. Modern Scientific Chemistry. Trans. by M.	_	٠,
M. Pattison Muir	*,	00
Latimer, L. H., Field, C. J., and Howell, J. W. Incandescent	_	•
Electric Lighting. (Science Series No. 57.)16mo,	•	50
Latta, M. N. Handbook of American Gas-Engineering Practice.	U	50
8vo.	*.	50
American Producer Gas Practice	_	90
Leask, A. R. Breakdowns at Sea		00
—— Triple and Quadruple Expansion Engines 12mo,		00
Refrigerating Machinery		00
Lecky, S. T. S. "Wrinkles" in Practical Navigation 8vo,	• **	00
Le Doux, M. Ice-Making Machines. (Science Series No. 46.)		
16mo,	0	50
Leeds, C. C. Mechanical Drawing for Trade Schools oblong, 4to,	_	
High School Edition		25
Machinery Trades Edition	*2	00
Lefévre, L. Architectural Pottery. Trans. by H. K. Bird and		
W. M. Binns4to,	*7	50
Lehner, S. Ink Manufacture. Trans. by A. Morris and H.		
Robson	*2	50
Lemstrom, S. Electricity in Agriculture and Horticulture.		
8vo,	*1	50
Le Van, W. B. Steam-Engine Indicator. (Science Series No.		
78.)16mo,	0	50
Lewes, V. B. Liquid and Gaseous Fuels. (Westminster Series.)		
8vo,	*2	00
Lieber, B. F. Lieber's Standard Telegraphic Code 8vo,	*10	00
—— Code. German Edition8vo,	*10	00
	*10	00
, French Edition	*10	00
—— Terminal Index8vo,	*2	50
	*15	00

Lieber, B. F. Handy Tables4to,	*2	50
Bankers and Stockbrokers' Code and Merchants and		
Shippers' Blank Tables8vo,	*15	00
100,000,000 Combination Code 8vo,	*15	00
Engineering Code8vo,	*10	00
Livermore, V. P., and Williams, J. How to Become a Com-		
petent Motorman	*1	00
Livingstone, R. Design and Construction of Commutators. 8vo.		25
Lobben, P. Machinists' and Draftsmen's Handbook 8vo,		ີ50ີ
Locke, A. G. and C. G. Manufacture of Sulphuric Acid 8vo,		00
Lockwood, T. D. Electricity, Magnetism, and Electro-teleg-		
raphy	2	50
— Electrical Measurement and the Galvanometer 12mo,		50
Lodge, O. J. Elementary Mechanics		50
		00
Lord, R. T. Decorative and Fancy Fabrics		50
Loring, A. E. A Handbook of the Electromagnetic Telegraph.	3	50
<b>-</b>	_	
(Science Series No. 39)		50
Loewenstein, L. C., and Crissey, C. P. Centrifugal Pumps. (In		
Lucke, C. E. Gas Engine Design	-3	00
Power Plants: their Design, Efficiency, and Power Costs.		\
2 vols(In Prepa	rano	n.)
Power Plant Papers. Form I. The Steam Power Plant	_	
paper,		50
Lunge, G. Coal-tar Ammonia. Two Volumes8vo,	*15	00
Manufacture of Sulphuric Acid and Alkali. Three Volumes		
Vol. I. Sulphuric Acid. In two parts	*15	00
Vol. II. Salt Cake, Hydrochloric Acid and Leblanc Soda.		
In two parts	*15	
Vol. III. Ammonia Soda	*15	
— Technical Chemists' Handbook12mo, leather,	*3	50
—— Technical Methods of Chemical Analysis. Trans. by C. A.		
Keane. In collaboration with the corps of specialists.		
Vol. I. In two parts8vo,	*15	00
Vols. II and III(In Prepa	ratio	n.)
Lupton, A., Parr, G. D. A., and Perkin, H. Electricity as Applied		
to Mining8vo,	*4	50
Tuguer I M Minerals in Pock Sections 800		Ĕ٥

Macewen, H. A. Food Inspection8vo,	*2	50
Mackenzie, N. F. Notes on Irrigation Works 8vo,		50
Mackie, J. How to Make a Woolen Mill Pay8vo,	*2	00
Mackrow, C. Naval Architect's and Shipbuilder's Pocket-		,
book16mo, leather,	5	00
Maguire, Capt. E. The Attack and Defense of Coast Fortifica-	Ŭ	
tions8vo,	2	50
Maguire, Wm. R. Domestic Sanitary Drainage and Plumbing		
8vo,	4	00
Mallet, A. Compound Engines. Trans. by R. R. Buel.	•	-
(Science Series No. 10.)		
Mansfield, A. N. Electro-magnets. (Science Series No. 64)		
16mo,		50
Marks, E. C. R. Construction of Cranes and Lifting Machinery	·	30
12mo,	*-	50
—— Construction and Working of Pumps		50
— Manufacture of Iron and Steel Tubes		00
		00
— Mechanical Engineering Materials		
—— Inventions, Patents and Designs12mo,		50
Markham, E. R. The American Steel Worker12mo,		00
		50
Marlow, T. G. Drying Machinery and Practice8vo,	_	00
Marsh, C. F. Concise Treatise on Reinforced Concrete8vo,		50
Marsh, C. F., and Dunn, W. Reinforced Concrete4to,	⁻5	00
Manual of Reinforced Concrete and Concrete Block Con-		
struction	*2	50
Marshall, W.J., and Sankey, H. R. Gas Engines. (Westminster		
Series.)	*2	00
Telephony12mo,	*1	00
Matheson, D. Australian Saw-Miller's Log and Timber Ready	7	
Reckoner 12mo, leather,		50
Mathot, R. E. Internal Combustion Engines8vo,	*6	00
Maurice, W. Electric Blasting Apparatus and Explosives 8vo,	*3	
Shot Firer's Guide8vo,	*1	50
Maxwell, J. C. Matter and Motion. (Science Series No. 36.)		
16mo,	0	50
Maxwell, W. H., and Brown, J. T. Encyclopedia of Municipal		
and Sanitary Engineering4to,	*10	00
Mayer, A. M. Lecture Notes on Physics8vo,	2	90

McCullough, R. S. Mechanical Theory of Heat8vo,	3	50
McIntosh, J. G. Technology of Sugar8vo,	*4	50
—— Industrial Alcohol	*3	00
- Manufacture of Varnishes and Kindred Industries. Three	-	
Volumes. 8vo.		
Vol. I. Oil Crushing, Refining and Boiling	*3	50
Vol. II. Varnish Materials and Oil Varnish Making	*4	_
Vol. III(In Prepar		
McKnight, J. D., and Brown, A. W. Marine Multitubular		
Boilers	*1	50
McMaster, J. B. Bridge and Tunnel Centres. (Science Series		
No. 20.)	0	50
McMechen, F. L. Tests for Ores, Minerals and Metals12mo,	*I	-
McNeill, B. McNeill's Code8vo,	*6	00
McPherson, J. A. Water-works Distribution 8vo,	2	
Melick, C. W. Dairy Laboratory Guide	*1	_
Merck, E. Chemical Reagents; Their Purity and Tests8vo,	*1	_
Merritt, Wm. H. Field Testing for Gold and Silver . 16mo, leather,	I	_
Meyer, J. G. A., and Pecker, C. G. Mechanical Drawing and		_
Machine Design4to,	5	00
Michell, S. Mine Drainage8vo,	10	
Mierzinski, S. Waterproofing of Fabrics. Trans. by A. Morris		
and H. Robson8vo,	*2	50
Miller, E. H. Quantitative Analysis for Mining Engineers . 8vo,	*1	_
Miller, G. A. Determinants. (Science Series No. 105.). 16mo,		•
Milroy, M. E. W. Home Lace-making	*I	00
Minifie, W. Mechanical Drawing8vo,	*4	00
Mitchell, C. A., and Prideaux, R. M. Fibres Used in Textile and	•	
Allied Industries		
Modern Meteorology	1	50
Monckton, C. C. F. Radiotelegraphy. (Westminster Series.)		
8vo,	*2	00
Monteverde, R. D. Vest Pocket Glossary of English-Spanish,		
Spanish-English Technical Terms 64mo, leather,	*I	00
Moore, E. C. S. New Tables for the Complete Solution of		
Ganguillet and Kutter's Formula8vo,	*5	00
Moreing, C. A., and Neal, T. New General and Mining Tele-	-	
graph Code8vo,	*5	00
Morgan, A. P. Wireless Telegraph Construction for Amateurs.	-	
12m0.	*T	K۵

•		
Moses, A. J. The Characters of Crystals8vo,	*2	2 00
Moses, A. J., and Parsons, C. I. Elements of Mineralogy. 8vo,	*2	2 50
Moss, S. A. Elements of Gas Engine Design. (Science		
Series.)16mo,	0	50
The Lay-out of Corliss Valve Gears. (Science Series). 16mo,	o	50
Mullin, J. P. Modern Moulding and Pattern-making 12mo,	2	50
Munby, A. E. Chemistry and Physics of Building Materials.		
(Westminster Series.)8vo,	*2	00
Murphy, J. G. Practical Mining16mo,	1	00
Murray, J. A. Soils and Manures. (Westminster Series.) 8vo,	*2	00
Naquet, A. Legal Chemistry12mo,	2	00
Nasmith, J. The Student's Cotton Spinning8vo,	3	00
Neilson, R. M. Aeroplane Patents8vo,	*2	00
Nerz, F. Searchlights. Trans. by C. Rodgers8vo,	*3	00
Neuberger, H., and Noalhat, H. Technology of Petroleum.		
Trans. by J. G. McIntosh8vo,	*10	00
Newall, J. W. Drawing, Sizing and Cutting Bevel-gears.		
8vo,	I	50
Newlands, J. Carpenters and Joiners' Assistant		
folio, half mor.,	15	00
Nicol, G. Ship Construction and Calculations8vo,	*4	50
Nipher, F. E. Theory of Magnetic Measurements 12mo,	I	00
Nisbet, H. Grammar of Textile Design8vo,	*3	00
Nolan, H. The Telescope. (Science Series No. 51.)16mo,	0	50
Noll, A. How to Wire Buildings	1	50
Nugent, E. Treatise on Optics	I.	50
O'Connor, H. The Gas Engineer's Pocketbook 12mo, leather,	3	50
—— Petrol Air Gas12mo,	*o '	75
Ohm, G. S., and Lockwood, T. D. Galvanic Circuit. Trans. by		
William Francis. (Science Series No. 102.) 16mo,	0	50
Olsen, J. C. Text book of Quantitative Chemical Analysis 8vo,	*4 (	00
Olsson, A. Motor Control, in Turret Turning and Gun Elevating.		
(U. S. Navy Electrical Series, No. 1.) 12mo, paper,	*o :	50
Oudin, M. A. Standard Polyphase Apparatus and Systems 8vo,	*3 (	ю.
Palaz, A. Industrial Photometry. Trans. by G. W. Patterson,		
Ir8vo.	*4 (	00

Pamely, C. Colliery Manager's Handbook8vo,	*10	00
Parr, G. D. A. Electrical Engineering Measuring Instruments.		
, 8vo.	*3	50
Parry, E. J. Chemistry of Essential Oils and Artificial Per-	·	•
fumes	*<	00
Parry, E. J., and Coste, J. H. Chemistry of Pigments 8vo,	_	50
Parry, L. A. Risk and Dangers of Various Occupations 8vo,		00
Parshall, H. F., and Hobart, H. M. Armature Windings 4to,	_	. 20
- Electric Railway Engineering	*10	-
Parshall, H. F., and Parry, E. Electrical Equipment of Tram-		
ways(In	Pres	ss.)
Parsons, S. J. Malleable Cast Iron		50
Passmore, A. C. Technical Terms Used in Architecture 8vo,		50
Patterson, D. The Color Printing of Carpet Yarns8vo,	_	50
Color Matching on Textiles8vo,	-	00
— The Science of Color Mixing8vo,	-	00
Patton, H. B. Lecture Notes on Crystallography 8vo,	_	25
Paulding, C. P. Condensation of Steam in Covered and Bare		Ī
Pipes8vo,	*2	00
Transmission of Heat through Cold-storage Insulation		
12mo,	*1	00
Peirce, B. System of Analytic Mechanics4to,		00
Pendred, V. The Railway Locomotive. (Westminster Series.)		
8vo.	*2	00
Perkin, F. M. Practical Methods of Inorganic Chemistry.		
12m0,	*1	00
Perrigo, O. E. Change Gear Devices8vo,		00
Perrine, F. A. C. Conductors for Electrical Distribution 8vo,		50
Perry, J. Applied Mechanics8vo,	-	50
Petit, G. White Lead and Zinc White Paints8vo,		50
Petit, R. How to Build an Aeroplane. Trans. by T. O'B.		•
Hubbard, and J. H. Ledeboer8vo,	*1	50
Pettit, Lieut. J. S. Graphic Processes. (Science Series No. 76.)		•
16mo,	o	50
Philbrick, P. H. Beams and Girders. (Science Series No. 88.)		
16mo,		
Phillips, J. Engineering Chemistry8vo,	*4	50
— Gold Assaying		50
— Dangerous Goods8vo,		50
	•	~

Phin, J. Seven Follies of Science	* I	25
Household Pests, and How to Get Rid of Them		
. 8vo (In Prepa	rati	on.)
Pickworth, C. N. The Indicator Handbook. Two Volumes		
12mo, each,	I	50
Logarithms for Beginners		50
— The Slide Rule 12mo,		00
Plane Table, The8vo,	2	00
Plattner's Manual of Blowpipe Analysis. Eighth Edition, re-		
vised. Trans. by H. B. Cornwall8vo,	*4	00
Plympton, G. W. The Aneroid Barometer. (Science Series.)		
16mo,	0	50
How to become an Engineer. (Science Series No. 100.)		
. r6mo,	0	50
Van Nostrand's Table Book. (Science Series No. 104.)		-
16mo,	0	50
Pochet, M. L. Steam Injectors. Translated from the French.		
(Science Series No. 29.)16mo,	0	50
Pocket Logarithms to Four Places. (Science Series.)16mo,	0	50
leather,	1	00
Pope, F. L. Modern Practice of the Electric Telegraph 8vo,	I	50
Popplewell, W. C. Elementary Treatise on Heat and Heat	•	
Engines12mo,	*3	00
Prevention of Smoke8vo,	*3	50
Strength of Minerals8vo,	*1	75
Potter, T. Concrete8vo,	*3	00
Practical Compounding of Oils, Tallow and Grease 8vo,	*3	50
Practical Iron Founding	1	50
Pray, T., Jr. Twenty Years with the Indicator8vo,	2	50
Steam Tables and Engine Constant8vo,	2 (	00
— Calorimeter Tables	í I (	00
Preece, W. H. Electric Lamps(In	Press	3.) `
Prelini, C. Earth and Rock Excavation8vo,	*3 (	00
—— Dredges and Dredging8vo (In	Press	3.)
Graphical Determination of Earth Slopes8vo,	*2 (	00
Tunneling	3 (	00
Prescott, A. B. Organic Analysis8vo,	5, 0	90
Prescott, A. B., and Johnson, O. C. Quanitative Chemical		
A neltrois Syo	*	<b>*</b> ^

D A D . 10 W . D 0 D 1 D 1 1 0 W		
Prescott, A. B., and Sullivan, E. C. First Book in Qualitative	_	
Chemistry12m0,	*1	50
Pritchard, O. G. The Manufacture of Electric-light Carbons.		
8vo, paper,	*0	60
Prost, E. Chemical Analysis of Fuels, Ores, Metals. Trans.		
by J. C. Smith 8vo,	*4	50
Pullen, W. W. F. Application of Graphic Methods to the Design		
of Structures12mo,	*2	50
Injectors: Theory, Construction and Working12mo,	*1	50
Pulsifer, W. H. Notes for a History of Lead8vo,	4	00
Purchase, W. R. Masonry12mo,	*3	00
Putsch, A. Gas and Coal-dust Firing8vo,	· <b>*</b> 3	00
Pynchon, T. R. Introduction to Chemical Physics8vo,	3	00
Rafter, G. W. Mechanics of Ventilation. (Science Series No.		
33.)16mo,	0	50
— Potable Water. (Science Series No. 103.)16mo,		50
Treatment of Septic Sewage. (Science Series.).	-	0-
16mo,	0	50
Rafter, G. W., and Baker, M. N. Sewage Disposal in the United	·	50
States4to,	*6	00
Raikes, H. P. Sewage Disposal Works		00
Railway Shop Up-to-Date		00
Ramp, H. M. Foundry Practice(In		
Randall, P. M. Quartz Operator's Handbook12mo,		00.) 00
Randau, P. Enamels and Enamelling		
Rankine, W. J. M. Applied Mechanics	•	00
	_	00
— Civil Engineering		50
— Machinery and Millwork8vo,	-	00
The Steam-engine and Other Prime Movers	-	00
— Useful Rules and Tables	4	00
Rankine, W. J. M., and Bamber, E. F. A Mechanical Text-		
book8vo,	3	50
Raphael, F. C. Localization of Faults in Electric Light and		
Power Mains8vo,	_	00
Rathbone, R. L. B. Simple Jewellery8vo,	*2	00
Rateau, A. Flow of Steam through Nozzles and Orifices.		
Trans. by H. B. Brydon8vo,	*1	50
Rausenberger, F. The Theory of the Recoil of Guns 8vo.	*4	50

Rautenstrauch, W. Notes on the Elements of Machine Design, 8vo, boards,	*1	. <b>5</b> 0
Rautenstrauch, W., and Williams, J. T. Machine Drafting and Empirical Design.		
Part I. Machine Drafting8vo,	* I	25
Part II. Empirical Design(In Prepa		
Raymond, E. B. Alternating Current Engineering12mo,		50
Rayner, H. Silk Throwing and Waste Silk Spinning 8vo,		50
Recipes for the Color, Paint, Varnish, Oil, Soap and Drysaltery		0-
Trades8vo,	*3	50
Recipes for Flint Glass Making	_	50
Redwood, B. Petroleum. (Science Series No. 92.)16mo,		50
Reed's Engineers' Handbook8vo,		00
Key to the Nineteenth Edition of Reed's Engineers' Hand-	•	
book8vo,	*2	00
— Useful Hints to Sea-going Engineers	-	50
Marine Boilers		00
Reinhardt, C. W. Lettering for Draftsmen, Engineers, and	_	-
Studentsoblong 4to, boards,		00
— The Technic of Mechanical Drafting oblong 4to, boards,	*1	
Reiser, F. Hardening and Tempering of Steel. Trans. by A.	•	•
Morris and H. Robson12mo,	**	50
Reiser, N. Faults in the Manufacture of Woolen Goods. Trans.	-	50
by A. Morris and H. Robson8vo,	*2	
— Spinning and Weaving Calculations	*5	-
Renwick, W. G. Marble and Marble Working8vo,	-	00
Reynolds, O., and Idell, F. E. Triple Expansion Engines.	5	00
(Science Series No. 99.)	*1	50
Rhead, G. W. British Pottery Marks		
	3	00
Rice, J. M., and Johnson, W. W. A New Method of Obtaining		42
the Differential of Functions		50
Richardson, J. The Modern Steam Engine8vo,	*3	-
Richardson, S. S. Magnetism and Electricity12mo,		00
Rideal, S. Glue and Glue Testing8vo,	*4	
Rings, F. Concrete in Theory and Practice	*2	
Ripper, W. Course of Instruction in Machine Drawing. folio,	+0	00
Roberts, F. C. Figure of the Earth. (Science Series No. 79.)		
16mo,	0	50

•		
Roberts, J., Jr. Laboratory Work in Electrical Engineering.		
8vo,	*2	00
Robertson, L. S. Water-tube Boilers8vo,	3	00
Robinson, J. B. Architectural Composition8vo,	*2	50
Robinson, S. W. Practical Treatise on the Teeth of Wheels.		
(Science Series No. 24.)	0	50
Railroad Economics. (Science Series No. 59.)16mo,	0	50
Wrought Iron Bridge Members. (Science Series No. 60.)		
16mo,	0	50
Robson, J. H. Machine Drawing and Sketching(In	Pres	88.)
Roebling, J. A. Long and Short Span Railway Bridges. folio,	25	00
Rogers, A. A Laboratory Guide of Industrial Chemistry . 12mo,	*I	50
Rogers, A., and Aubert, A. B. Industrial Chemistry(In	Pres	38.)
Rogers, F. Magnetism of Iron Vessels. (Science Series No. 30.)		
16то,	0	50
Rollins, W. Notes on X-Light8vo,		00
Rose, J. The Pattern-makers' Assistant8vo,	2	50
- Key to Engines and Engine-running12mo,	2	50
Rose, T K. The Precious Metals. (Westminster Series.).		
8vo,	*2	00
Rosenhain, W. Glass Manufacture. (Westminster Series.) 8vo,	*2	οó
Ross, W. A. Blowpipe in Chemistry and Metallurgy12mo,	*2	00
Rossiter, J. T. Steam Engines. (Westminster Series.)		
8vo (In	Pres	ss.)
Pumps and Pumping Machinery. (Westminster Series.)		•
8vo (In .	Pres	88.)
Roth. Physical Chemistry8vo,	*2	00
Rouillion, L. The Economics of Manual Training8vo,	2	00
Rowan, F. J. Practical Physics of the Modern Steam-boiler		
8vo.	7	50
Rowan, F. J., and Idell, F. E. Boiler Incrustation and Corro-	٠	•
sion. (Science Series No. 27.)	0	50
Roxburgh, W. General Foundry Practice8vo,	*3	-
Ruhmer, E. Wireless Telephony. Trans. by J. Erskine-		
Murray8vo,	*3	50
Russell, A. Theory of Electric Cables and Networks8vo,	*3	-
	_	
Sabine, R. History and Progress of the Electric Telegraph. 12mo,	I	25
Saeltzer, A. Treatise on Acoustics		00

Salomons, D. Electric Light Installations. 12mo.		
Vol. I. The Management of Accumulators	2	50
Vol. II. Apparatus	2	25
Vol. III. Applications	I	50
Sanford, P. G. Nitro-explosives8vo,	*4	00
Saunders, C. H. Handbook of Practical Mechanics 16mo,	I	00
leather,	1	25
Saunnier, C. Watchmaker's Handbook12mo,	3	00
Sayers, H. M. Brakes for Tram Cars8vo,	*1	25
Scheele, C. W. Chemical Essays8vo,	*2	00
Schellen, H. Magneto-electric and Dynamo-electric Machines		
8vo,	5	00
Scherer, R. Casein. Trans. by C. Salter 8vo,	*3	00
Schmall, C. N. First Course in Analytic Geometry, Plane and		
Solid12mo, half leather,	*1	75
Schmall, C. N., and Schack, S. M. Elements of Plane Geometry		
12m0,	*1	25
Schmeer, L. Flow of Water8vo,		00
Schumann, F. A Manual of Heating and Ventilation.		
12mo, leather,	1	50
Schwartz, E. H. L. Causal Geology8vo,		50
Schweizer, V., Distillation of Resins8vo,		50
Scott, W. W. Qualitative Chemical Analysis. A Laboratory	•	•
Manual8vo	* *I	50
Scribner, J. M. Engineers' and Mechanics' Companion.		•
16mo, leather,	I	50
Searle, G. M. "Sumners' Method." Condensed and Improved.		
(Science Series No. 124.)8vo,	6	00
Seaton, A. E. Manual of Marine Engineering8vo,	6	00
Seaton, A. E., and Rounthwaite, H. M. Pocket-book of Marine	٠.	
Engineering16mo, leather,	· з	00
Seeligmann, T., Torrilhon, G. L., and Falconnet, H. India	Ū	
Rubber and Gutta Percha. Trans. by J. G. McIntosh		
8vo,	*5	00
Seidell, A. Solubilities of Inorganic and Organic Substances	•	
8vo,	*3	00
Sellew, W. H. Steel Rails4to (In	_	
Senter, G. Outlines of Physical Chemistry12mo,		50
Sever. G. F. Electric Engineering Experiments 8vo. hoards.		-

Same C. P. and Tammand P. Jahanatam and Pastom Tests		
Sever, G. F., and Townsend, F. Laboratory and Factory Tests	*-	50
in Electrical Engineering		_
Sewall, C. H. Wireless Telegraphy8vo,		00
Lessons in Telegraphy		00
Sewell, T. Elements of Electrical Engineering 8vo,	_	00
—— The Construction of Dynamos8vo,	-	00
Sexton, A. H. Fuel and Refractory Materials 12mo,		50
—— Chemistry of the Materials of Engineering 12mo,		50
—— Alloys (Non-Ferrous) 8vo,	_	00
—— The Metallurgy of Iron and Steel8vo,		50
Seymour, A. Practical Lithography8vo,	*2	50
— Modern Printing Inks8vo,	*2	00
Shaw, H. S. H. Mechanical Integrators. (Science Series No.		٠
83.)16mo,	0	50
Shaw, P. E. Course of Practical Magnetism and Electricity.		
8vo,	*1	00
Shaw, S. History of the Staffordshire Potteries8vo,	*3	00
Chemistry of Compounds Used in Porcelain Manufacture	•	
8vo,	*5	00
Sheldon, S., and Hausmann, E. Direct Current Machines.	Ŭ	
8vo.	*2	50
— Alternating-current Machines		50
Sherer, R. Casein. Trans. by C. Salter8vo,		00
Sherriff, F. F. Oil Merchants' Manual	_	50
Shields, J. E. Notes on Engineering Construction12mo,	-	50
Shock, W. H. Steam Boilers4to, half mor.,	,	90
Shreve, S. H. Strength of Bridges and Roofs	_	
Shunk, W. F. The Field Engineer		50
	2	50
Simmons, W. H., and Appleton, H. A. Handbook of Soap	٠.	
Manufacture8vo,	_	00
Simms, F. W. The Principles and Practice of Leveling8vo,		50
—— Practical Tunneling		.50
Simpson, G. The Naval Constructor 12mo, mor.,	*5	00
Sinclair, A. Development of the Locomotive Engine.		
8vo, half leather,	5	00
Sindall, R. W. Manufacture of Paper. (Westminster Series.)		
8 <b>v</b> o,	*2	00
Sloane, T. O'C. Elementary Electrical Calculations 12mo,	*2	00
Smith, C. A. M. Handbook of Testing, Vol. I. Materials (In.	Pres	RR.)

Smith, C. F. Practical Alternating Currents and Testing8vo,	*2 50
Practical Testing of Dynamos and Motors8vo,	*2 00
Smith, F. E. Handbook of General Instruction for Mechanics.	
12mo,	I 50
Smith, I. W. The Theory of Deflections and of Latitudes and	
Departures16mo, mor.,	3 00
Smith, J. C. Manufacture of Paint8vo,	*3 00
Smith, W. Chemistry of Hat Manufacturing12mo,	*3 00
Snell, A. T. Electric Motive Power8vo,	*4 00
Snow, W. G. Pocketbook of Steam Heating and Ventilation	
	Press.)
Snow, W. G., and Nolan, T. Ventilation of Buildings. (Science	•
Series No. 5.)	O 50
Soddy, F. Radioactivity8vo,	*3 00
Solomon, M. Electric Lamps. (Westminster Series.)8vo,	*2 00
Sothern, J. W. The Marine Steam Turbine8vo,	*5 00
Soxhlet, D. H. Dyeing and Staining Marble. Trans. by A.	
Morris and H. Robson 8vo,	*2 50
Spang, H. W. A Practical Treatise on Lightning Protection	•
12mo,	I 00
Spangenburg, L. Fatigue of Metals. Translated by S. H.	
Shreve. (Science Series No. 23.)	0 50
Specht, G. J., Hardy, A. S., McMaster, J. B., and Walling. Topo-	•
graphical Surveying. (Science Series No. 72.) 16mo,	0 50
Spevers, C. L. Text-book of Physical Chemistry8vo.	*2 25
Stahl, A. W. Transmission of Power. (Science Series No. 28.)	
16mo,	
Stahl, A. W., and Woods, A. T. Elementary Mechanism . 12mo,	*2 00
Staley, C., and Pierson, G. S. The Separate System of Sewerage.	
8vo,	*3 00
Standage, H. C. Leatherworkers' Manual8vo,	*3 50
— Sealing Waxes, Wafers, and Other Adhesives 8vo,	*2 00
- Agglutinants of all Kinds for all Purposes 12mo,	*3 50
Stansbie, J. H. Iron and Steel. (Westminster Series.) 8vo,	*2 00
Stevens, H. P. Paper Mill Chemist 16mo,	*2 50
Stevenson, J. L. Blast-Furnace Calculations12mo, leather,	*2 00
Stewart, A. Modern Polyphase Machinery12mo,	*2 00
Stewart, G. Modern Steam Traps12mo,	*I 25
Stiles, A. Tables for Field Engineers	I 00

 $\mathbf{r}$ 

G.111		
Stillman, P. Steam-engine Indicator12mo,		00
Stodola, A. Steam Turbines. Trans. by L. C. Loewenstein. 8vo,	_	00
Stone, H. The Timbers of Commerce8vo,	3	50
Stone, Gen. R. New Roads and Road Laws12mo,	I	00
Stopes, M. Ancient Plants8vo,		00
The Study of Plant Life8vo (In	Pres	88.)
Sudborough, J. J., and James, T. C. Practical Organic Chem-		
istry12m0,	*2	00
Suffling, E. R. Treatise on the Art of Glass Painting 8vo,	*3	50
Swan, K. Patents, Designs and Trade Marks. (Westminster		
Series.)8vo,	*2	00
Sweet, S. H. Special Report on Coal8vo,	3	00
Swinburne, J., Wordingham, C. H., and Martin, T. C. Electric	Ū	
Currents. (Science Series No. 109.)16mo,	0	50
Swoope, C. W. Practical Lessons in Electricity12mo,		00
• • •		
Tailfer, L. Bleaching Linen and Cotton Yarn and Fabrics 8vo,	*5	00
Tate, J. S. Surcharged and Different Forms of Retaining-walls.	Ŭ	
(Science Series No. 7.)16mo,		
Templeton, W. Practical Mechanic's Workshop Companion.		
12mo, mor.,	2	00
Terry, H. L. India Rubber and its Manufacture. (Westminster	_	•
Series.)	*2	^^
Thiess, J. B., and Joy, G. A. Toll Telephone Practice (In Prepa		
Thom, C., and Jones, W. H. Telegraphic Connections.		,
oblong 12mo,		50
Thomas, C. W. Paper-makers' Handbook(In		
Thompson, A. B. Oil Fields of Russia	*7	
—— Petroleum Mining and Oil Field Development 8vo,		-
Thompson, E. P. How to Make Inventions	*5	
Thompson, S. P. Dynamo Electric Machines. (Science Series	U	50
	_	
No. 75.)	O	50
Thompson, W. P. Handbook of Patent Law of All Countries	_	
I6mo,		50
Thornley, T. Cotton Combing Machines8vo,	*3	00
Cotton Spinning8vo,	•	
First Year	*1	
Second Year	*2	
Third Year	*2	50

Thurso, J. W. Modern Turbine Practice	*4 0
94.)	0 50
Tinney, W. H. Gold-mining Machinery8vo.	*3 00
Titherley, A. W. Laboratory Course of Organic Chemistry	
8vo.	*2 00
Toch, M. Chemistry and Technology of Mixed Paints	
8vo,	*3 00
_	Press.)
Todd, J., and Whall, W. B. Practical Seamanship 8vo,	*7 50
Tonge, J. Coal. (Westminster Series.)8vo,	*2 00
Townsend, F. Alternating Current Engineering8vo, boards,	*o 75
Townsend, J. Ionization of Gases by Collision8vo,	*I 25
Transactions of the American Institute of Chemical Engineers.	
8vo,	
Vol. I. 1908	*6 <b>oo</b>
Vol. II. 1909	*6 <b>o</b> o
Traverse Tables. (Science Series No. 115.)16mo,	0 50
mor.,	I 00
Trinks, W., and Housum, C. Shaft Governors. (Science	
Series No. 122.)	0 50
Trowbridge, W. P. Turbine Wheels. (Science Series No. 44.)	
16mo,	0 50
Tucker, J. H. A Manual of Sugar Analysis8vo,	3 50
Tumlirz, O. Potential. Trans. by D. Robertson12mo,	1 25
Tunner, P. A. Treatise on Roll-turning. Trans. by J. B.	
Pearse	10 00
Turbayne, A. A. Alphabets and Numerals4to,	2 00
Turnbull, Jr., J., and Robinson, S. W. A Treatise on the Com-	
pound Steam-engine. (Science Series No. 8.)16mo,	
Turrill, S. M. Elementary Course in Perspective 12mo,	*I 25
Underhill, C. R. Solenoids, Electromagnets and Electromag-	
netic Windings12mo,	*2 00
Urquhart, J. W. Electric Light Fitting	2 00
— Electro-plating	2 00
— Electrotyping	` 2 00
— Electric Ship Lighting	3 00
Universal Telegraph Cipher Code	I 00

#### d. van nostrand company's short-title catalog 35

Vacher, F. Food Inspector's Handbook	*2 50
Van Nostrand's Chemical Annual. Second issue 1909 12mo,	*2 50
Year Book of Mechanical Engineering Data. First issue	
1910(In	Press.)
Van Wagenen, T. F. Manual of Hydraulic Mining 16mo,	I 00
Vega, Baron, Von Logarithmic Tables8vo, half mor.,	2 50
Villon, A. M. Practical Treatise on the Leather Industry.	
	*10 00
Vincent, C. Ammonia and its Compounds. Trans. by M. J.	
Salter8vo,	<b>*</b> 2 00
Volk, C. Haulage and Winding Appliances8vo,	*4 00
Von Georgiovics, G. Chemical Technology of Textile Fibres.	
Trans. by C. Salter	*4 50
Chemistry of Dyestuffs. Trans. by C. Salter8vo,	*4 50
Vose, G. L. Graphic Method for Solving Certain Questions in	
Arithmetic and Algebra. (Science Series No. 16.)16mo,	0 50
Wabner, R. Ventilation in Mines. Trans. by C. Salter 8vo,	*4 50
Wade, E. J. Secondary Batteries8vo,	*4 00
Wadsworth, C. Primary Battery Ignition 12mo (In	Press.)
Wagner, E. Preserving Fruits, Vegetables, and Meat12mo,	*2 50
Walker, F. Aerial Navigation 8vo,	
— Dynamo Building. (Science Series No. 98.)16mo,	0 50
—— Electric Lighting for Marine Engineers 8vo,	2 00
Walker, S. F. Steam Boilers, Engines and Turbines 8vo,	3 00
— Refrigeration, Heating and Ventilation on Shipboard.	
12m0,	*2 00
—— Electricity in Mining 8vo,	*3 50
Walker, W. H. Screw Propulsion8vo,	0 75
Wallis-Tayler, A. J. Bearings and Lubrication8vo,	*1 50
— Modern Cycles 8vo,	4 00
—— Motor Cars	1 80
— Motor Vehicles for Business Purposes8vo,	3 50
— Pocket Book of Refrigeration and Ice Making12mo,	I 50
Refrigerating and Ice-making Machinery8vo,	3 00
Refrigeration and Cold Storage	*4 50
—— Sugar Machinery 12mo,	*2 00
Wanklyn, J. A. Treatise on the Examination of Milk 12mo,	1 00
— Water Analysis 12mo,	2 00

Wansbrough, W. D. The A B C of the Differential Calculus	
. 12m0,	*I 5
Slide Valves	*2 0
Ward, J. H. Steam for the Million	IO
Waring, Jr., G. E. Sanitary Conditions. (Science Series No. 31.)	
16mo.	0 5
Sewerage and Land Drainage	*6 o
Modern Methods of Sewage Disposal	2 0
How to Drain a House	I 2
Warren, F. D. Handbook on Reinforced Concrete12mo,	*2 50
Watkins, A. Photography. (Westminster Series)8vo (In	
Watson, E. P. Small Engines and Boilers	
Watt, A. Electro-plating and Electro-refining of Metals	I 25
Electro-metallurgy 12mo,	*4 50
— The Art of Paper Making	I 00
	*3 00
The Art of Soap-making	3 00
— Leather Manufacture8vo,	*4 00
Weale, J. Dictionary of Terms used in Architecture12mo,	2 50
Weale's Scientific and Technical Series. (Complete list sent on	
application.)	
Weather and Weather Instruments12m0,	I 00
paper,	0 50
Webb, H. L. Guide to the Testing of Insulated Wires and	
Cables12mo,	I 00
Webber, W. H. Y. Town Gas. (Westminster Series.)8vo,	*2 00
Weekes, R. W. The Design of Alternate Current Transformers	
12mo,	I 00
Weisbach, J. A Manual of Theoretical Mechanics8vo,	*6 oo
sheep,	*7 50
Weisbach, J., and Herrmann, G. Mechanics of Air Machinery	
8vo,	*3 75
Weston, E. B. Loss of Head Due to Friction of Water in Pipes	
12m0,	*I 50
Weymouth, F. M. Drum Armatures and Commutators.	•
8vo,	*3 00
Wheeler, J. B. Art of War	1 75
—— Field Fortifications 12mo,	I 75
Whipple, S. An Elementary and Practical Treatise on Bridge	- 13
Duttitue	

Whithard, P. Illuminating and Missal Painting12mo, Wilcox, R. M. Cantilever Bridges. (Science Series No. 25.)	I	50
16mo.	0	50
Wilkinson, H. D. Submarine Cable Laying and Repairing.	_	5-
8vo.	*6	00
· •	.0	00
Williams, A. D., Jr., and Hutchinson, R. W. The Steam Turbine.	<b>D</b>	\
•	Pres	
Williamson, R. S. On the Use of the Barometer4to,	15	
—— Practical Tables in Meteorology and Hypsometery4to,		50
Willson, F. N. Theoretical and Practical Graphics4to,	*4	00
Wimperis, H. E. Internal Combustion Engine8vo,	*3	00
Winchell, N. H., and A. N. Elements of Optical Mineralogy.		
8vo,	*3	50
Winkler, C., and Lunge, G. Handbook of Technical Gas-Analy-		
sis	4	00
Winslow, A. Stadia Surveying. (Science Series No. 77.).	•	
16mo,	, ,	50
Wisser, Lieut. J. P. Explosive Materials. (Science Series No.	_	0-
70.)16mo,	•	50
Modern Gun Cotton. (Science Series No. 89.) 16mo,		50
Wood, De V. Luminiferous Aether. (Science Series No. 85.)	U	30
•	_	
16mo,	0	50
Woodbury, D. V. Elements of Stability in the Well-propor-		
tioned Arch8vo, half mor.,	4	00
Worden, E. C. The Nitrocellulose Industry. Two Volumes.	_	
8vo (In		
Wright, A. C. Analysis of Oils and Allied Substances8vo,		50
Simple Method for Testing Painter's Materials8vo,	*2	50
Wright, H. E. Handy Book for Brewers8vo,	*5	00
Wright, F. W. Design of a Condensing Plant12mo,	*1	50
Wright, T. W. Elements of Mechanics8vo,	*2	50
Wright, T. W., and Hayford, J. F. Adjustment of Observations		
8vo,	*3	00
	•	
Young, J. E. Electrical Testing for Telegraph Engineers8vo,	*4	00
Zahner, R. Transmission of Power. (Science Series No. 40.)		
Iomo,		
•	*.	
Zeidler, J., and Lustgarten, J. Electric Arc Lamps8vo,	₹2	00

Zeune	r, A.	Technical	Thermodynamics.	Trans.	by J. F.		
	Klei	n. Two V	olumes		8vo,	*8	00
Zimm	er, G. I	. Mechan	ical Handling of Ma	terial	4to,	*ro	00
Zipeer,	, j. T	extile Raw l	Materials. Trans. b	y C. Salte	r8vo,	*5	00
Zur N	ledden,	F. Engin	eering Workshop M	lachines	and Proc-		
	6686	a. Trans.	by I. A. Davenport .		8vo.	*2	00

Books sent postpaid to any address on receipt of price.

Descriptive circulars and complete catalogs may be had on application.

## FOSTIR

#### **ELECTRICAL**

#### **ENGINEER'S**

man and Enterpol o

#### **POCKET BOOK**

инна 1600 pp. 1128 ffl. 716 Tobles ga

Foster's Pocketbook is the busy engineer's first assistant. When he wants really reliable information, he turns to Foster to get it for him, and he gets it. Foster places before him data on the latest and best practice in electrical

engineering with a minimum expenditure of time and effort.

The material in Foster is logically arranged and is indexed in a full table of contents and a voluminous index that alone covers sixty pages, and to this is added a set of patent thumb index tabs that make reference to any section of the book practically instantaneous. The index is most thorough and reliable. It points right to the spot where the information sought is.

If you ever need information on electricity, you ought to have a copy of the new Fifth Edition, completely revised and enlarged, with four-fifths of the old matter replaced by new, up-to-date material, and containing all the excellent features named above, with 1,636 pages, 1,128 illustrations and 718 tables. The price is \$5.00. ORDER A COPY NOW

# CKETB

By ALBERT I. FRYE, M. Am. Soc. C. E.

A COMPREHENSIVE treatment of Civil Engineering, in seventy sections, comprising about 1,400 pages; with 500 tables and 1,000 illustrations. Also a complete glossary of engineering terms.

Each main subject receives economic consideration and analysis, and is reinforced with excerpts from, and references to the most important cost and other data in our leading technical publications—including hundreds of illustrations of up-to-date engineering structures and details.

"How to design engineering structures economically" has received careful study in the preparation of this work. The fundamental principles laid down can be followed readily by every young engineer, and carried to practical completion in the finished structure.

Most of the tables are new and have been prepared regardless of time and expense. Many of them have been arranged in a novel manner which will appeal especially to engineers; and all have been carefully checked and rechecked to eliminate the possibility of errors.

Graphical methods have been illustrated freely throughout the work,

in connection with analytical solutions.

Methods and cost of work have received special attention; and, thus, there is crowded into the 1,400 pages of text, in small type and in pocketbook form, a vast amount of engineering data, formulas and tables, that will be found absolutely indispensable to every engineer, contractor and student.

D. VAN NOSTRAND COMPANY, Publishers 23 Murray and 27 Warren Sts., **NEW YORK** 

• . . . 

\* .

